Cite as: S. Zhao *et al.*, *Science*
10.1126/science.aat2299 (2018).

Enantiodivergent Pd-catalyzed C–C bond formation enabled through ligand parameterization

Shibin Zhao,^{1,2*} Tobias Gensch,^{3*} Benjamin Murray,^{1,2} Zachary L. Niemeyer,³ Matthew S. Sigman,^{3†} Mark R. Biscoe^{1,2†}

¹Department of Chemistry, The City College of New York, 160 Convent Avenue, New York, NY 10031, USA. ²Ph.D. Program in Chemistry, The Graduate Center of the City University of New York, 365 Fifth Avenue, New York, NY 10016, USA. ³Department of Chemistry, University of Utah, 315 South 1400 East, Salt Lake City, UT 84112, USA.

*These authors contributed equally to this work. †Corresponding author. Email: sigman@chem.utah.edu (M.S.S.); mbiscoe@ccny.cuny.edu (M.R.B.)

Despite the enormous potential for the use of stereospecific cross-coupling reactions to rationally manipulate the three-dimensional structure of organic molecules, the factors that control the transfer of stereochemistry in these reactions remain poorly understood. Herein we report a mechanistic and synthetic investigation into the use of enantioenriched alkylboron nucleophiles in stereospecific Pd-catalyzed Suzuki cross-coupling reactions. By developing a suite of molecular descriptors of phosphine ligands, we could apply predictive statistical models to select or design distinct ligands that respectively promoted stereoinvertive and stereoretentive cross-coupling reactions. Stereodefined branched structures were thereby accessed through the predictable manipulation of absolute stereochemistry, and a general model for the mechanism of alkylboron transmetallation was proposed.

Palladium-catalyzed cross-coupling reactions have revolutionized the construction of $C(sp^2)$ – $C(sp^2)$ bonds. Among these cross-coupling processes, the Suzuki–Miyaura reaction has found particularly broad application due to its extensive reaction scope, as well as the stability, availability, and low toxicity of organoboron reagents (*1*). The 2010 Nobel Prize in chemistry was awarded, in part, to recognize the transformative impact of the Suzuki cross-coupling reaction on chemical synthesis. However, although $C(sp^2)$ – $C(sp^2)$ bond construction is now considered routine using the Suzuki reaction, extension of this process to the formation of $C(sp^3)$ – $C(sp^2)$ bonds using alkylboron nucleophiles remains a significant challenge. Of particular interest, a variant using secondary alkylboron nucleophiles with predictable and controllable stereospecificity would establish a powerful synthetic strategy to access molecular geometries with precise three-dimensional control, expanding the exceptional capabilities of the Suzuki reaction (Fig. 1A).

Many efforts have focused on the use of enantioenriched secondary alkylboron nucleophiles in Suzuki cross-coupling reactions (*2–4*). Significant limitations remain due to slow transmetallation of the highly covalent and sterically congested $C(sp^3)$ –B bond in these reagents, as well as the propensity of the resulting Pd-alkyl species to undergo β -hydride elimination/reinsertion sequences, which can result in isomerization of the alkyl group and racemization of the stereocenter. To circumvent prohibitively slow transmetallation, as well as competing β -hydride elimination/reinsertion pathways, most stereospecific Suzuki reactions have required the use of secondary alkylboron nucleophiles that are

electronically activated via inclusion of a $C(sp^2)$ α -carbon, an α -heteroatom, and/or a strongly coordinating β -carbonyl group (*5–17*). In addition, alkylboron nucleophiles can undergo transmetallation via either stereoretentive or stereoinvertive pathways depending on the nature of the substrate, catalyst, and/or reaction conditions. In many cases, the factors controlling the dominant mechanism of transmetallation are not understood (Fig. 1B). Thus, a predictive stereochemical model for transmetallation of alkylboron reagents remains elusive.

Recently, we reported a stereospecific Pd-catalyzed cross-coupling reaction using unactivated secondary alkylboron nucleophiles (*18*). With P^tBu_3 as a supporting ligand, enantioenriched arylation products were obtained with transmetallation proceeding primarily via a stereoinvertive mechanism. Whereas several enlightening mechanistic studies have recently been conducted on the transmetallation of arylboron nucleophiles (*19–23*), these studies have not addressed the transmetallation of alkylboron nucleophiles in $C(sp^3)$ – $C(sp^2)$ bond-forming processes (*24, 25*). Thus, unactivated alkylboron nucleophiles constitute an attractive starting point from which to investigate the reaction parameters most influential to the mechanism of alkylboron transmetallation. This mechanistic work should simultaneously facilitate the development of new synthetic methods to rationally incorporate/manipulate stereocenters via cross-coupling strategies. To this end, we report a study using predictive statistical models (*26, 27*) to relate phosphine ligand properties to stereochemical outcomes obtained from Pd-catalyzed Suzuki reactions of unactivated enantioenriched secondary alkylboron

nucleophiles and aryl electrophiles. With statistical models that rely on a next generation set of molecular descriptors, we achieved a stereoretentive Pd-catalyzed cross-coupling reaction of such nucleophiles. Furthermore, we have identified an improved ligand for the stereoinvertive variant, thus enabling an entirely ligand-controlled enantiodivergent process from a single-enantiomer organoboron nucleophile (28). Our statistical models also provide compelling evidence that each transmetallation pathway is intimately tied to specific electronic properties of the supporting ligand, which serves as a predictive guide to the mechanism of alkylboron transmetalation to palladium.

Initial investigations using electronically differentiated aryl chlorides with enantioenriched $^8\text{BuBF}_3\text{K}$ revealed a trend correlating diminished stereofidelity with the use of more electron-deficient coupling partners (Fig. 1C, I). This observation suggested that subtle electronic effects could influence the mechanism of transmetallation and the resulting stereochemical outcome. Additionally, when the phosphine ligand was varied in an initial screen with a common aryl chloride electrophile, a considerable change in the reaction outcome from stereoinversion to stereoretention was found (Fig. 1C, II). No obvious correlation was observed between these results and the steric properties (solid angle) of the ligand. Taken together, these outcomes were difficult to interpret and inspired the use of ligand parameterization tools to provide a platform for both predictive ligand performance and mechanistic interrogation.

An expanded inventory of common phosphines with varied properties was evaluated in the Suzuki reaction of enantioenriched $^8\text{BuBF}_3\text{K}$ and ethyl 4-chlorobenzoate. This dataset was then subjected to correlation analysis of phosphine structural features with the stereochemical outcomes as well as the ratios of branched:linear products in these reactions. We devised a workflow and universal parameter set to describe the catalyst properties from the phosphine itself (29–32). The workflow was initiated by performing a molecular mechanics (MM) conformational search to reveal representative low energy conformers. (Fig. 2A). Next, geometry optimization of the conformers using DFT was followed by parameter collection. Subsequently, four descriptor subsets were defined to capture the conformational dynamics of the ligands by including the mathematical extreme descriptor values (minimum and maximum), the lowest energy conformer values, and the Boltzmann weighted averages. We viewed the unique treatment of representative conformers as a crucial means of describing ensemble properties such as chemical shift while also probing structural flexibility during catalysis.

The final step in the workflow involved the analysis of both the stereofidelity and the branched:linear product ratio. These two readouts presumably describe two stages of the reaction mechanism (Fig. 1B): i) the competing stereoretentive

and stereoinvertive transmetallation mechanisms that determine the final stereochemistry of the cross-coupling product and ii) the competitive β -hydride elimination/isomerization sequences that follow transmetallation. A correlation of the branched:linear ratio with the final enantiopurity of the product reveals that β -hydride elimination is responsible for both racemization and isomerization to the linear side product. Furthermore, a modest trend is observed relating the minimum width B_1 of the phosphine ligand to the branched:linear ratio (Fig. 2B). This is consistent with reports of large ligands facilitating reductive elimination over β -hydride elimination (33), and suggests the use of a parameterization approach to take into account the conformational flexibility of ligands.

Since the inherent selectivity of the transmetallation mechanism is masked by deleterious racemization as a consequence of β -hydride elimination, only ligands providing high selectivity were further investigated (>30% ee, Fig. 2B). The molecular electrostatic potential minimum in the phosphorus lone pair region (V_{min}) has been shown to correlate with the classical Tolman electronic parameter (34). Thus, V_{min} serves as an easily computable measure for the overall ligand electronics. A correlation between enantioselectivity and V_{min} was observed within the abridged dataset indicating that electronic properties of the ligand determine the mechanism of transmetallation. Specifically, electron-rich trialkylphosphines promoted stereoinvertive reactions, whereas the electron-poorer triarylphosphines provided modest selectivity for stereoretention. Use of the bulky, electron-rich ligand PAd_3 (**9**), which was recently reported by Carrow (35), resulted in a particularly large preference for the stereoinvertive outcome. Based on these data, we hypothesized and virtually evaluated ligands for improved stereoretentive outcomes with the following features: i) large ligand bulk to prevent β -hydride elimination and racemization, and ii) electron-deficient aryl substituents at phosphorus to promote the stereoretentive mechanism and to accelerate reductive elimination (Fig. 2B). Among the proposed ligands was a set of biaryl phosphines (**11–15**), as pioneered by Buchwald (36), featuring various electron-deficient aryl groups at phosphorus. Gratifyingly, ligands **11** and **14** promote the alkyl Suzuki cross-coupling reaction with significantly enhanced selectivity (up to 90% ee) and minimal alkyl isomerization. Thus, parameterization-driven optimization facilitated development of a stereoretentive Suzuki reaction involving unactivated alkylboron nucleophiles. When considered alongside the introduction of **9** to achieve stereoinvertive couplings, predictable control of the absolute sense of enantioselectivity (retention or inversion) can be engendered by simply selecting the appropriate ligand.

Our stereochemical investigations of secondary alkylboron transmetallation in the Suzuki reaction suggested that both enantiomers of a cross-coupling product could be

selectively accessed through use of a single enantioenriched alkylboron reagent with the proper selection of the phosphine ligand. The scope of this process is depicted in Fig. 3. Using enantioenriched, unactivated alkyltrifluoroborate nucleophiles, ligand-controlled stereoselectivity was broadly achieved in cross-coupling reactions with aryl electrophiles. Strongly π -accepting ligands bis- $\text{CF}_3\text{PhSPhos}$ (**11**) and bis- $\text{CF}_3\text{PhXPhos}$ (**14**), which emerged from our parameterization-guided optimization, preferentially promote the stereoretentive pathway, while strongly σ -donating ligand PAd_3 (**9**) preferentially promotes the stereoinvertive pathway. Because electron-poor palladium catalysts commonly undergo slow oxidative addition with aryl chlorides, we also evaluated aryl bromide and triflate electrophiles in reactions involving **11** and **14**. A particular highlight of this protocol is the uniformity of the conditions used for both the stereoinvertive and stereoretentive reactions: each operates in a toluene/water mixture as solvent, with a carbonate base, and no additional additives. Both reaction variants tolerated the use of electron-rich and electron-deficient aryl electrophiles, as well as an aryl electrophile bearing an *ortho*-substituent. High stereofidelity was achieved for all of these reactions, including those involving alkylboron nucleophiles bearing thiophenyl and phenoxide substituents. Use of an alkylboron nucleophile containing a larger substituent (replacing methyl with ethyl) at the stereogenic center was also well tolerated (**16i**). In the absence of a methyl substituent, the stereoinvertive variant shows modestly reduced selectivity that may be improved by reducing the reaction temperature. This was partially anticipated due to the greater sensitivity of the metal fragment to steric congestion at carbon in the stereoinvertive $\text{S}_{\text{E}}2$ transmetallation mechanism. Diastereomeric products **17a** and **17b** could be generated from a single alkylboron diastereomer (**37**) using **14** and PAd_3 , respectively (Fig. 3B). In these reactions, replacement of ligand **14** with PAd_3 resulted in a change in diastereoselectivity from 30:1 to 1:5, a 3.6 kcal/mol free energy of activation difference dependent only on the ligand identity. No erosion of specificity was observed for electron-deficient aryl substrates in stereoinvertive Suzuki reactions using PAd_3 , in contrast to analogous reactions using P^tBu_3 . Furyl and thiophenyl electrophiles are also compatible with our system (Fig. 3C). As an additional mechanistic probe, *trans*-2-methylcyclopentyltrifluoroborate was subjected to the stereodivergent reaction conditions. Because *trans*-2-methylcyclopentyltrifluoroborate is sterically impeded from undergoing stereoinvertive transmetallation, only the stereoretentive process using **14** should be mechanistically viable. Indeed, we observed that use of ligand **14** smoothly generates **18** with stereoretention, while use of PAd_3 results in low alkylboron conversion.

To further probe the origin of the ligand-dependent enantiodivergent process, we interrogated the mechanism of

transmetallation using the parameterization strategy described above (Fig. 4). To accomplish this, phenyl-substituted substrate **20** was selected due to enhanced performance and thus a greater output range. A singular aryl chloride electrophile was chosen for this analysis to avoid potential attenuation of the ligand effects by the influence of different counterions (e.g., bromide or triflate). Additionally, 24 ligands were tested, excluding smaller ligands to reduce the complexity associated with β -hydride elimination. Multivariate linear regression revealed that most of the outputs can be expressed in two readily interpretable terms that discriminate the transmetallation pathways: the average energy of the P–C antibonding orbitals $E_{\sigma^*(\text{P-C})}$, representative of π -back bonding, and the energy of the lone pair orbital of phosphorus $E_{\text{LP}(\text{P})}$, a measure of the ligand's σ -donation capability (Fig. 4B). This outcome suggests that the stereoinvertive pathway is dependent on strong σ -donation from the ligand, which may stabilize a two-coordinate, cationic palladium complex. Conversely, the stereoretentive pathway is enhanced by π -back bonding, which may stabilize the coordination of a π -donor ligand X (presumably OH^-) to Pd. Including two steric descriptors such as B_1 and L improves the model fit by treating the competitive β -hydride elimination that occurs using smaller ligands and decreases the observed specificity. This becomes evident when the four smallest ligands in this dataset are removed from the analysis, which results in an excellent correlation using just the two electronic descriptors with the experimentally observed stereochemical outcomes (Fig. 4D). Multivariate regression analysis thereby provides compelling evidence for the electronic factors favoring each transmetallation mechanism and thus a guideline for future developments in stereospecific cross-coupling reactions.

REFERENCES AND NOTES

1. J. C. H. Lee, D. G. Hall, "State-of-the-art in metal-catalyzed cross-coupling reactions of organoboron compounds with organic electrophiles," in *Metal-Catalyzed Cross-Coupling Reactions and More*, A. de Meijere, S. Bräse, M. Oestreich, Eds. (Wiley-VCH, ed. 3, 2014), vol. 1, pp. 65–132.
2. C.-Y. Wang, J. Derosaa, M. R. Biscoe, Configurationally stable, enantioenriched organometallic nucleophiles in stereospecific Pd-catalyzed cross-coupling reactions: An alternative approach to asymmetric synthesis. *Chem. Sci.* **6**, 5105–5113 (2015). [doi:10.1039/C5SC01710F](https://doi.org/10.1039/C5SC01710F) [Medline](#)
3. C. Sandford, V. K. Aggarwal, Stereospecific functionalizations and transformations of secondary and tertiary boronic esters. *Chem. Commun. (Camb.)* **53**, 5481–5494 (2017). [doi:10.1039/C7CC01254C](https://doi.org/10.1039/C7CC01254C) [Medline](#)
4. J. P. G. Rygus, C. M. Crudden, Enantiospecific and iterative Suzuki-Miyaura cross-couplings. *J. Am. Chem. Soc.* **139**, 18124–18137 (2017). [doi:10.1021/jacs.7b08326](https://doi.org/10.1021/jacs.7b08326) [Medline](#)
5. D. Imao, B. W. Glasspoole, V. S. Laberge, C. M. Crudden, Cross coupling reactions of chiral secondary organoboronic esters with retention of configuration. *J. Am. Chem. Soc.* **131**, 5024–5025 (2009). [doi:10.1021/ja8094075](https://doi.org/10.1021/ja8094075) [Medline](#)
6. D. L. Sandrock, L. Jean-Gérard, C. Y. Chen, S. D. Dreher, G. A. Molander, Stereospecific cross-coupling of secondary alkyl β -trifluoroboratoamides. *J. Am. Chem. Soc.* **132**, 17108–17110 (2010). [doi:10.1021/ja108949w](https://doi.org/10.1021/ja108949w) [Medline](#)
7. T. Ohmura, T. Awano, M. Suginoe, Stereospecific Suzuki-Miyaura coupling of chiral α -(acylamino)benzylboronic esters with inversion of configuration. *J. Am. Chem. Soc.* **132**, 13191–13193 (2010). [doi:10.1021/ja106632j](https://doi.org/10.1021/ja106632j) [Medline](#)

8. T. Awano, T. Ohmura, M. Suginome, Inversion or retention? Effects of acidic additives on the stereochemical course in enantiospecific Suzuki-Miyaura coupling of α -(acetylamino)benzylboronic esters. *J. Am. Chem. Soc.* **133**, 20738–20741 (2011). [doi:10.1021/ja210025q Medline](#)
9. J. C. H. Lee, R. McDonald, D. G. Hall, Enantioselective preparation and chemoselective cross-coupling of 1,1-diboron compounds. *Nat. Chem.* **3**, 894–899 (2011). [doi:10.1038/nchem.1150 Medline](#)
10. B. M. Partridge, L. Chausset-Boissarie, M. Burns, A. P. Pulis, V. K. Aggarwal, Enantioselective synthesis and cross-coupling of tertiary propargylic boronic esters using lithiation-borylation of propargylic carbamates. *Angew. Chem. Int. Ed.* **51**, 11795–11799 (2012). [doi:10.1002/anie.201203198 Medline](#)
11. G. A. Molander, S. R. Wisniewski, Stereospecific cross-coupling of secondary organotrifluoroborates: Potassium 1-(benzyloxy)alkyltrifluoroborates. *J. Am. Chem. Soc.* **134**, 16856–16868 (2012). [doi:10.1021/ja307861n Medline](#)
12. S. C. Matthew, B. W. Glasspoole, P. Eisenberger, C. M. Crudden, Synthesis of enantiomerically enriched triarylmethanes by enantiospecific Suzuki-Miyaura cross-coupling reactions. *J. Am. Chem. Soc.* **136**, 5828–5831 (2014). [doi:10.1021/ja42159g Medline](#)
13. C. Sun, B. Potter, J. P. Morken, A catalytic enantiotopic-group-selective Suzuki reaction for the construction of chiral organoboronates. *J. Am. Chem. Soc.* **136**, 6534–6537 (2014). [doi:10.1021/ja500029w Medline](#)
14. T. P. Blaisdell, J. P. Morken, Hydroxyl-directed cross-coupling: A scalable synthesis of debromohamigeran E and other targets of interest. *J. Am. Chem. Soc.* **137**, 8712–8715 (2015). [doi:10.1021/jacs.5b05477 Medline](#)
15. S. Miyamura, M. Araki, T. Suzuki, J. Yamaguchi, K. Itami, Stereodivergent synthesis of arylcyclopropylamines by sequential C–H borylation and Suzuki-Miyaura coupling. *Angew. Chem. Int. Ed.* **54**, 846–851 (2015). [doi:10.1002/anie.201409186 Medline](#)
16. Y. Lou, P. Cao, T. Jia, Y. Zhang, M. Wang, J. Liao, Copper-catalyzed enantioselective 1,6-boration of para-quinone methides and efficient transformation of gem-diarylmethine boronates to triarylmethanes. *Angew. Chem. Int. Ed.* **127**, 12302–12306 (2015). [doi:10.1002/ange.201505926](#)
17. G. L. Hoang, J. M. Takacs, Enantioselective γ -borylation of unsaturated amides and stereoretentive Suzuki-Miyaura cross-coupling. *Chem. Sci.* **8**, 4511–4516 (2017). [doi:10.1039/C7SC01093A Medline](#)
18. L. Li, S. Zhao, A. Joshi-Pangu, M. Diane, M. R. Biscoe, Stereospecific Pd-catalyzed cross-coupling reactions of secondary alkylboron nucleophiles and aryl chlorides. *J. Am. Chem. Soc.* **136**, 14027–14030 (2014). [doi:10.1021/ja508815w Medline](#)
19. C. Amatore, A. Jutand, G. Le Duc, Kinetic data for the transmetalation/reductive elimination in palladium-catalyzed Suzuki-Miyaura reactions: Unexpected triple role of hydroxide ions used as base. *Chemistry* **17**, 2492–2503 (2011). [doi:10.1002/chem.201001911 Medline](#)
20. B. P. Carrow, J. F. Hartwig, Distinguishing between pathways for transmetalation in Suzuki-Miyaura reactions. *J. Am. Chem. Soc.* **133**, 2116–2119 (2011). [doi:10.1021/ja1108326 Medline](#)
21. A. A. Thomas, S. E. Denmark, Pre-transmetalation intermediates in the Suzuki-Miyaura reaction revealed: The missing link. *Science* **352**, 329–332 (2016). [doi:10.1126/science.aad6981 Medline](#)
22. A. A. Thomas, H. Wang, A. F. Zahrt, S. E. Denmark, Structural, kinetic, and computational characterization of the elusive arylpalladium(II)boronate complexes in the Suzuki-Miyaura reaction. *J. Am. Chem. Soc.* **139**, 3805–3821 (2017). [doi:10.1021/jacs.6b13384 Medline](#)
23. A. A. Thomas, A. F. Zahrt, C. P. Delaney, S. E. Denmark, Elucidating the role of the boronic esters in the Suzuki-Miyaura reaction: Structural, kinetic, and computational investigations. *J. Am. Chem. Soc.* **140**, 4401–4416 (2018). [doi:10.1021/jacs.8b00400 Medline](#)
24. B. H. Ridgway, K. A. Woerpel, Transmetalation of alkylboranes to palladium in the Suzuki coupling reaction proceeds with retention of stereochemistry. *J. Org. Chem.* **63**, 458–460 (1998). [doi:10.1021/jo970803d Medline](#)
25. K. Matos, J. A. Soderquist, Alkylboranes in the Suzuki-Miyaura coupling: Stereochemical and mechanistic studies. *J. Org. Chem.* **63**, 461–470 (1998). [doi:10.1021/jo971681s Medline](#)
26. C. B. Santiago, J.-Y. Guo, M. S. Sigman, Predictive and mechanistic multivariate linear regression models for reaction development. *Chem. Sci.* **9**, 2398–2412 (2018). [doi:10.1039/C7SC04679K Medline](#)
27. M. S. Sigman, K. C. Harper, E. N. Bess, A. Milo, The development of multidimensional analysis tools for asymmetric catalysis and beyond. *Acc. Chem. Res.* **49**, 1292–1301 (2016). [doi:10.1021/acs.accounts.6b00194 Medline](#)
28. S. Krautwald, E. M. Carreira, Stereodivergence in asymmetric catalysis. *J. Am. Chem. Soc.* **139**, 5627–5639 (2017). [doi:10.1021/jacs.6b13340 Medline](#)
29. C. A. Tolman, Steric effects of phosphorus ligands in organometallic chemistry and homogeneous catalysis. *Chem. Rev.* **77**, 313–348 (1977). [doi:10.1021/cr60307a002](#)
30. J. Jover, N. Fey, J. N. Harvey, G. C. Lloyd-Jones, A. G. Orpen, G. J. J. Owen-Smith, P. Murray, D. R. J. Hose, R. Osborne, M. Purdie, Expansion of the ligand knowledge base for monodentate P-donor ligands (LKB-P). *Organometallics* **29**, 6245–6258 (2010). [doi:10.1021/om100648v](#)
31. Z. L. Niemeyer, A. Milo, D. P. Hickey, M. S. Sigman, Parameterization of phosphine ligands reveals mechanistic pathways and predicts reaction outcomes. *Nat. Chem.* **8**, 610–617 (2016). [doi:10.1038/nchem.2501 Medline](#)
32. K. Wu, A. G. Doyle, Parameterization of phosphine ligands demonstrates enhancement of nickel catalysis via remote steric effects. *Nat. Chem.* **9**, 779–784 (2017). [doi:10.1038/nchem.2741 Medline](#)
33. R. Jana, T. P. Pathak, M. S. Sigman, Advances in transition metal (Pd, Ni, Fe)-catalyzed cross-coupling reactions using alkyl-organometallics as reaction partners. *Chem. Rev.* **111**, 1417–1492 (2011). [doi:10.1021/cr100327p Medline](#)
34. C. H. Suresh, N. Koga, Quantifying the electronic effect of substituted phosphine ligands via molecular electrostatic potential. *Inorg. Chem.* **41**, 1573–1578 (2002). [doi:10.1021/ic0109400 Medline](#)
35. L. Chen, P. Ren, B. P. Carrow, Tri(1-adamantyl)phosphine: Expanding the boundary of electron-releasing character available to organophosphorus compounds. *J. Am. Chem. Soc.* **138**, 6392–6395 (2016). [doi:10.1021/jacs.6b03215 Medline](#)
36. D. S. Surry, S. L. Buchwald, Dialkylbiaryl phosphines in Pd-catalyzed amination: A user's guide. *Chem. Sci.* **2**, 27–50 (2011). [doi:10.1039/C0SC00331I Medline](#)
37. S. Balieu, G. E. Hallett, M. Burns, T. Bootwicha, J. Studley, V. K. Aggarwal, Toward ideality: The synthesis of (+)-kalkitoxin and (+)-hydroxyphthioceranic acid by assembly-line synthesis. *J. Am. Chem. Soc.* **137**, 4398–4403 (2015). [doi:10.1021/ja512875g Medline](#)
38. H. Li, L. Wang, Y. Zhang, J. Wang, Transition-metal-free synthesis of pinacol alkylboronates from tosylhydrazones. *Angew. Chem. Int. Ed.* **51**, 2943–2946 (2012). [doi:10.1002/anie.201108139 Medline](#)
39. V. Bagutski, A. Ros, V. K. Aggarwal, Improved method for the conversion of pinacolboronic esters into trifluoroborate salts: Facile synthesis of chiral secondary and tertiary trifluoroborates. *Tetrahedron* **65**, 9956–9960 (2009). [doi:10.1016/j.tet.2009.10.002](#)
40. G. A. Molander, L. N. Cavalcanti, Oxidation of organotrifluoroborates via oxone. *J. Org. Chem.* **76**, 623–630 (2011). [doi:10.1021/jo22208d Medline](#)
41. S. D. Dreher, P. G. Dormer, D. L. Sandrock, G. A. Molander, Efficient cross-coupling of secondary alkyltrifluoroborates with aryl chlorides—Reaction discovery using parallel microscale experimentation. *J. Am. Chem. Soc.* **130**, 9257–9259 (2008). [doi:10.1021/ja8031423 Medline](#)
42. W. F. Bailey, X. Jiang, Cyclization of (4-methoxy-5-hexenyl)lithium. *J. Org. Chem.* **59**, 6528–6533 (1994). [doi:10.1021/jo00101a010](#)
43. R. Larouche-Gauthier, T. G. Elford, V. K. Aggarwal, Ate complexes of secondary boronic esters as chiral organometallic-type nucleophiles for asymmetric synthesis. *J. Am. Chem. Soc.* **133**, 16794–16797 (2011). [doi:10.1021/ja2077813 Medline](#)
44. A. Bonet, M. Odachowski, D. Leonori, S. Essafi, V. K. Aggarwal, Enantiospecific sp^2 - sp^3 coupling of secondary and tertiary boronic esters. *Nat. Chem.* **6**, 584–589 (2014). [doi:10.1038/nchem.1971 Medline](#)
45. M. Burns, S. Essafi, J. R. Bame, S. P. Bull, M. P. Webster, S. Balieu, J. W. Dale, C. P. Butts, J. N. Harvey, V. K. Aggarwal, Assembly-line synthesis of organic molecules with tailored shapes. *Nature* **513**, 183–188 (2014). [doi:10.1038/nature13711 Medline](#)
46. T. E. Barder, S. D. Walker, J. R. Martinelli, S. L. Buchwald, Catalysts for Suzuki-Miyaura coupling processes: Scope and studies of the effect of ligand structure. *J. Am. Chem. Soc.* **127**, 4685–4696 (2005). [doi:10.1021/ja042491j Medline](#)
47. N. Hartmann, M. Niemeyer, Easy large-scale synthesis of sterically encumbered 2-iodobiphenyls and some derivatives thereof. *Syn. Comm.* **31**, 3839–3845 (2001).

- [doi:10.1081/SCC-100108234](https://doi.org/10.1081/SCC-100108234)
48. J. D. Hicks, A. M. Hyde, A. M. Cuezva, S. L. Buchwald, Pd-catalyzed *N*-arylation of secondary acyclic amides: Catalyst development, scope, and computational study. *J. Am. Chem. Soc.* **131**, 16720–16734 (2009). [doi:10.1021/ja9044357](https://doi.org/10.1021/ja9044357) [Medline](#)
 49. N. C. Bruno, M. T. Tudge, S. L. Buchwald, Design and preparation of new palladium precatalysts for C–C and C–N cross-coupling reactions. *Chem. Sci.* **4**, 916–920 (2013). [doi:10.1039/C2SC20903A](https://doi.org/10.1039/C2SC20903A) [Medline](#)
 50. N. C. Bruno, N. Niljianskul, S. L. Buchwald, *N*-substituted 2-aminobiphenylpalladium methanesulfonate precatalysts and their use in C–C and C–N cross-couplings. *J. Org. Chem.* **79**, 4161–4166 (2014). [doi:10.1021/jo500355k](https://doi.org/10.1021/jo500355k) [Medline](#)
 51. S. D. Friis, T. Skrydstrup, S. L. Buchwald, Mild Pd-catalyzed aminocarbonylation of (hetero)aryl bromides with a palladacycle precatalyst. *Org. Lett.* **16**, 4296–4299 (2014). [doi:10.1021/ol502014h](https://doi.org/10.1021/ol502014h) [Medline](#)
 52. L. Li, C. Y. Wang, R. Huang, M. R. Biscoe, Stereoretentive Pd-catalyzed Stille cross-coupling reactions of secondary alkyl azastannatranes and aryl halides. *Nat. Chem.* **5**, 607–612 (2013). [doi:10.1038/nchem.1652](https://doi.org/10.1038/nchem.1652) [Medline](#)
 53. S. A. Green, J. L. Matos, A. Yagi, R. A. Shenvi, Branch-selective hydroarylation: Iodoarene-olefin cross-coupling. *J. Am. Chem. Soc.* **138**, 12779–12782 (2016). [doi:10.1021/jacs.6b08507](https://doi.org/10.1021/jacs.6b08507) [Medline](#)
 54. W. M. Czaplak, M. Mayer, A. J. V. Wangelin, Direct cobalt-catalyzed cross-coupling between aryl and alkyl halides. *Synlett* **18**, 2931–2934 (2009).
 55. A. Wilsily, Y. Nguyen, E. Fillion, Hydrogenolysis of unstrained carbon-carbon σ bonds: Stereoselective entry into benzylic tertiary centers. *J. Am. Chem. Soc.* **131**, 15606–15607 (2009). [doi:10.1021/ja9076815](https://doi.org/10.1021/ja9076815) [Medline](#)
 56. J. E. Baldwin, S. Bonacorsi Jr., Stereochemistry of the thermal isomerizations of (1*R*, 2*R*)-1-((*E*)-styryl)-2-methylcyclopropane to 3-phenyl-4-methylcyclopentenes. *J. Am. Chem. Soc.* **115**, 10621–10627 (1993). [doi:10.1021/ja00076a021](https://doi.org/10.1021/ja00076a021)
 57. Schrödinger Release 2018-1: MacroModel (Schrödinger, LLC, New York, 2018).
 58. K. Ermanis, K. E. B. Parkes, T. Agback, J. M. Goodman, Expanding DP4: Application to drug compounds and automation. *Org. Biomol. Chem.* **14**, 3943–3949 (2016). [doi:10.1039/C6OB00015K](https://doi.org/10.1039/C6OB00015K) [Medline](#)
 59. K. Ermanis, J. M. Goodman, PyDP4 workflow integrating MacroModel/TINKER, Gaussian/NWChem and DP4 analysis, version 0.7 (2016); <https://github.com/KristapsF/PyDP4/>
 60. M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, X. Li, M. Caricato, A. V. Marenich, J. Bloino, B. G. Janesko, R. Gomperts, B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, D. Williams-Young, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J. A. Montgomery Jr., J. E. Peralta, F. Ogliaro, M. J. Bearpark, J. J. Heyd, E. N. Brothers, K. N. Kudin, V. N. Staroverov, T. A. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. P. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C. Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman, D. J. Fox, Gaussian 16, Revision A.03 (Gaussian, Inc., Wallingford, CT, 2016).
 61. C. Adamo, V. Barone, Toward reliable density functional methods without adjustable parameters: The PBE0 model. *J. Chem. Phys.* **110**, 6158–6170 (1999). [doi:10.1063/1.478522](https://doi.org/10.1063/1.478522)
 62. W. J. Hehre, R. Ditchfield, J. A. Pople, Self-consistent molecular orbital methods. XII. Further extensions of Gaussian-type basis sets for use in molecular orbital studies of organic molecules. *J. Chem. Phys.* **56**, 2257–2261 (1972). [doi:10.1063/1.1677527](https://doi.org/10.1063/1.1677527)
 63. P. C. Hariharan, J. A. Pople, The influence of polarization functions on molecular orbital hydrogenation energies. *Theor. Chim. Acta* **28**, 213–222 (1973). [doi:10.1007/BF00533485](https://doi.org/10.1007/BF00533485)
 64. T. Clark, J. Chandrasekhar, G. W. Spitznagel, P. V. R. Schleyer, Efficient diffuse function-augmented basis sets for anion calculations. III. The 3-21+G basis set for first-row elements, Li–F. *J. Comput. Chem.* **4**, 294–301 (1983). [doi:10.1002/jcc.540040303](https://doi.org/10.1002/jcc.540040303)
 65. R. Krishnan, J. S. Binkley, R. Seeger, J. A. Pople, Self-consistent molecular orbital methods. XX. A basis set for correlated wave functions. *J. Chem. Phys.* **72**, 650–654 (1980). [doi:10.1063/1.438955](https://doi.org/10.1063/1.438955)
 66. J. R. Cheeseman, G. W. Trucks, T. A. Keith, M. J. Frisch, A comparison of models for calculating nuclear magnetic resonance shielding tensors. *J. Chem. Phys.* **104**, 5497–5509 (1996). [doi:10.1063/1.471789](https://doi.org/10.1063/1.471789)
 67. Y. Zhao, D. G. Truhlar, The M06 suite of density functionals for main group thermochemistry, thermochemical kinetics, noncovalent interactions, excited states, and transition elements: Two new functionals and systematic testing of four M06-class functionals and 12 other function. *Theor. Chem. Acc.* **120**, 215–241 (2008). [doi:10.1007/s00214-007-0310-x](https://doi.org/10.1007/s00214-007-0310-x)
 68. F. Weigend, R. Ahlrichs, Balanced basis sets of split valence, triple zeta valence and quadruple zeta valence quality for H to Rn: Design and assessment of accuracy. *Phys. Chem. Chem. Phys.* **7**, 3297–3305 (2005). [doi:10.1039/b508541a](https://doi.org/10.1039/b508541a) [Medline](#)
 69. E. D. Glendening, A. E. Reed, J. E. Carpenter, F. Weinhold, *NBO Version 3.1* (University of Wisconsin–Madison, 1990).
 70. S. K. Latypov, F. M. Polyancev, D. G. Yakhvarov, O. G. Sinyashin, Quantum chemical calculations of ^{31}P NMR chemical shifts: Scopes and limitations. *Phys. Chem. Chem. Phys.* **17**, 6976–6987 (2015). [doi:10.1039/C5CP00240K](https://doi.org/10.1039/C5CP00240K) [Medline](#)
 71. H. Clavier, S. P. Nolan, Percent buried volume for phosphine and *N*-heterocyclic carbene ligands: Steric properties in organometallic chemistry. *Chem. Commun. (Camb.)* **46**, 841–861 (2010). [doi:10.1039/b922984a](https://doi.org/10.1039/b922984a) [Medline](#)
 72. L. Falivene, R. Credendino, A. Poater, A. Petta, L. Serra, R. Oliva, V. Scarano, L. Cavallo, SambVca 2. A web tool for analyzing catalytic pockets with topographic steric maps. *Organometallics* **35**, 2286–2293 (2016). [doi:10.1021/acs.organomet.6b00371](https://doi.org/10.1021/acs.organomet.6b00371)
 73. SambVca 2.0 server; www.molnac.unisa.it/OMtools/sambvca2.0/index.html.
 74. A. Verloop, in *Drug Design*, E. J. Ariens, Ed. (Academic Press, 1976), vol. 3, pp. 133–187.
 75. T. Piou, F. Romanov-Michailidis, M. Romanova-Michaelides, K. E. Jackson, N. Semakul, T. D. Taggart, B. S. Newell, C. D. Rithner, R. S. Paton, T. Rovis, correlating reactivity and selectivity to cyclopentadienyl ligand properties in Rh(III)-catalyzed C–H activation reactions: An experimental and computational study. *J. Am. Chem. Soc.* **139**, 1296–1310 (2017). [doi:10.1021/jacs.6b11670](https://doi.org/10.1021/jacs.6b11670) [Medline](#)
 76. K. Jackson, R. Paton, Sterimol.py (2017); <https://github.com/bobbypaton/Sterimol>.
 77. I. A. Guzei, M. Wendt, An improved method for the computation of ligand steric effects based on solid angles. *Dalton Trans.* **2006**, 3991–3999 (2006). [doi:10.1039/b605102b](https://doi.org/10.1039/b605102b) [Medline](#)
 78. I. A. Guzei, M. Wendt, Program Solid-G (University of Wisconsin–Madison, 2004).
 79. MATLAB R2017a (Mathworks, Inc., Natick, MA, 2017).
 80. J.-Y. Guo, Y. Minko, C. B. Santiago, M. S. Sigman, Developing comprehensive computational parameter sets to describe the performance of pyridine-oxazoline and related ligands. *ACS Catal.* **7**, 4144–4151 (2017). [doi:10.1021/acscatal.7b00739](https://doi.org/10.1021/acscatal.7b00739)

ACKNOWLEDGMENTS

We acknowledge R. Kinthada for contributions to the diastereoselectivity studies. We thank G. Ralph for assistance with chiral HPLC analysis. We thank Prof. Luigi Cavallo for providing us with the command line tool for computing percent buried volume. **Funding:** we are grateful to the National Institutes of Health (grant SC1GM110010 to M.R.B.), the National Science Foundation (grant CHE-1665189 to M.R.B. and grants CHE-1361296 and CHE-1763436 to M.S.S.), and the Leopoldina Fellowship Programme of the German National Academy of Sciences Leopoldina (LPDS 2017-18 to T.G.). The support and resources from the Center for High Performance Computing (CHPC) at the University of Utah are gratefully acknowledged. Further computational resources were provided by the Extreme Science and Engineering Discovery Environment (XSEDE), which is supported by the NSF (ACI-1548562) and provided through allocation TG-CHE180003. **Author contributions:** S.Z. and B.M. performed all synthetic experiments and isolated all products. T.G. computed all ligand parameters and performed multivariate analyses. Z.L.N. performed initial multivariate analyses. M.R.B., M.S.S., and T.G. wrote the manuscript. M.R.B. and M.S.S. directed the project. **Competing interests:** The authors declare no competing interests. **Data and materials availability:** All additional data are in the supplementary materials.

SUPPLEMENTARY MATERIALS

www.sciencemag.org/cgi/content/full/science.aat2299/DC1

Materials and Methods

Supplementary Text

Fig. S1

Tables S1 to S6

NMR Spectra

References (38–80)

18 May 2018; accepted 6 September 2018

Published online 20 September 2018

10.1126/science.aat2299

Fig. 1. Reaction development. (A) Enantiodivergent Suzuki reactions of secondary alkylboron nucleophiles. (B) General mechanism. (C) Initial investigation of substrate and ligand influences on stereoselectivity. +% ee = net retention, -% ee = net inversion.

Fig. 2. Phosphine parameterization. (A) Workflow of parameter generation and statistical modeling. (B) Application of phosphine parameterization to ligand optimization of the reaction shown in Fig. 1C with Z = CO₂Et. b:l = branched to linear ratio. +% ee = net retention, -% ee = net inversion.

Fig. 3. Stereodivergent Pd-catalyzed cross-coupling reactions using enantioenriched alkylboron nucleophiles. The general reaction scheme is shown at the top. (A to C) Isolated yields are shown for stereoretentive and stereoinvertive cross-coupling reactions (A), diastereoselective cross-coupling reactions (B), and additional individual reactions (C). % es = % ee (final product) / % ee (starting material). [†] 44% yield, 84% ee (86% es) when run at 60 °C.

Fig. 4. Mechanistic investigation by multidimensional regression modeling. (A) Data from the arylation of **20** was used. [PdL] is the precatalyst as shown in Fig. 3 with varying ligands. (B) Regression model containing all 24 ligands of this data set. $E_{\sigma^*(P-C)_{avg}}^{Boltz}$: Boltzmann-weighted average across the conformers of the average energies of the three P-C σ^* antibonding orbitals in each phosphine. $E_{LP(P)}^{Boltz}$: Boltzmann-weighted average of the energy of the phosphorus lone pair orbital. Sterimol B_1^{Boltz} is the least width and Sterimol $L_{minconf}$ is the length of the lowest-energy conformer as seen from opposite the P substituents. Red points in the diagram: validation data (EV) not used in the model training. LOO = leave-one-out cross validation score. K-fold: average threefold cross-validation score. (C) Illustration and interpretation of the model terms. (D) Regression model after removing the four smallest ligands in this data set to exclude the influence of competitive β -hydride elimination on the data.