Cite as: M. Agostini *et al.*, *Science*
10.1126/science.aav8613 (2019).

Probing Majorana neutrinos with double- β decay

M. Agostini¹, A. M. Bakalyarov², M. Balata³, I. Barabanov⁴, L. Baudis⁵, C. Bauer⁶, E. Bellotti^{7,8}, S. Belogurov^{4,9}, A. Bettini^{10,11}, L. Bezrukov⁴, D. Borowicz¹², V. Brudanin¹², R. Brugnera^{10,11}, A. Caldwell¹³, C. Cattadori⁸, A. Chernogorov⁹, T. Comellato¹, V. D'Andrea¹⁴, E. V. Demidova⁹, N. DiMarco³, A. Domula¹⁵, E. Doroshkevich⁴, V. Egorov^{12*}, R. Falkenstein¹⁶, M. Fomina¹², A. Gangapshev⁴, A. Garfagnini^{10,11}, M. Giordano¹⁴, P. Grabmayr¹⁶, V. Gurentsov⁴, K. Gusev^{1,2,12}, J. Hakenmüller⁶, A. Hegai¹⁶, M. Heisel⁶, S. Hemmer¹¹, R. Hiller⁵, W. Hofmann⁶, M. Hult¹⁷, L. V. Inzhechik⁴, J. Janicskó Csáthy¹, J. Jochum¹⁶, M. Junker³, V. Kazalov⁴, Y. Kermaïdic⁶, T. Kihm⁶, I. V. Kirpichnikov⁹, A. Kirsch⁶, A. Kish⁵, A. Klimenko^{6,12}, R. Kneißl¹³, K. T. Knöpfle^{6†}, O. Kochetov¹², V. N. Kornoukhov^{4,9}, P. Krause¹, V. V. Kuzminov⁴, M. Laubenstein³, A. Lazzaro¹, M. Lindner⁶, I. Lippi¹¹, A. Lubashevskiy¹², B. Lubsandorzhiev⁴, G. Lutter¹⁷, C. Macolino³, B. Majorovits¹³, W. Maneschg⁶, M. Miloradovic⁵, R. Mingazheva⁵, M. Misiaszek¹⁸, P. Moseev⁴, I. Nemchenok¹², K. Panas¹⁸, L. Pandola¹⁹, K. Pelczar³, L. Pertoldi^{10,11}, P. Piseri²⁰, A. Pullia²⁰, C. Ransom⁵, S. Riboldi²⁰, N. Rumyantseva^{2,12}, C. Sada^{10,11}, E. Sala¹³, F. Salamida¹⁴, C. Schmitt¹⁶, B. Schneider¹⁵, S. Schönert¹, A.-K. Schütz¹⁶, O. Schulz¹³, M. Schwarz¹, B. Schwingenheuer⁶, O. Selivanenko⁴, E. Shevchik¹², M. Shirchenko¹², H. Singen⁶, A. Smolnikov^{6,12}, L. Stanco¹¹, D. Stukov², L. Vanhoefer¹³, A. A. Vasenko⁹, A. Veresnikova⁴, K. von Sturm^{10,11}, V. Wagner⁶, A. Wegmann⁶, T. Wester¹⁵, C. Wiesinger¹, M. Wojcik¹⁸, E. Yanovich⁴, I. Zhitnikov¹², S. V. Zhukov², D. Zinatulina¹², A. Zschocke¹⁶, A. J. Zsigmond¹³, K. Zuber¹⁵, G. Zuzel¹⁸, GERDA Collaboration

¹Physik Department, Technische Universität München, D-85748 Munich, Germany. ²National Research Centre "Kurchatov Institute," Moscow 123182, Russia. ³INFN Laboratori Nazionali del Gran Sasso and Gran Sasso Science Institute, I-67100 Assergi, Italy. ⁴Institute for Nuclear Research of the Russian Academy of Sciences, Moscow 117312, Russia. ⁵Physik Institut der Universität Zürich, CH-8057 Zurich, Switzerland. ⁶Max-Planck-Institut für Kernphysik, D-69117 Heidelberg, Germany. ⁷Dipartimento di Fisica, Università Milano Bicocca, I-20126 Milan, Italy. ⁸INFN Milano Bicocca, I-20126 Milan, Italy. ⁹Institute for Theoretical and Experimental Physics, Moscow 117259, Russia. ¹⁰Dipartimento di Fisica e Astronomia dell'Università di Padova, I-35121 Padua, Italy. ¹¹INFN Padova, I-35131 Padua, Italy. ¹²Joint Institute for Nuclear Research, Dubna 141980, Russia. ¹³Max-Planck-Institut für Physik, D-80805 Munich, Germany. ¹⁴INFN Laboratori Nazionali del Gran Sasso and Università degli Studi dell'Aquila, I-67100 L'Aquila, Italy. ¹⁵Institut für Kern- und Teilchenphysik, Technische Universität Dresden, D-01069 Dresden, Germany. ¹⁶Physikalisches Institut, Eberhard Karls Universität Tübingen, D-72076 Tübingen, Germany. ¹⁷European Commission, JRC-Geel, B-2440 Geel, Belgium. ¹⁸Institute of Physics, Jagiellonian University, Cracow 40-348, Poland. ¹⁹INFN Laboratori Nazionali del Sud, I-95123 Catania, Italy. ²⁰Dipartimento di Fisica, Università degli Studi di Milano e INFN Milano, I-20133 Milan, Italy.

*Deceased.

†Corresponding author. Email: gerda-eb@mpi-hd.mpg.de

A discovery that neutrinos are Majorana fermions would have profound implications for particle physics and cosmology. The Majorana character of neutrinos would make possible the neutrinoless double- β ($0\nu\beta\beta$) decay, a matter-creating process without the balancing emission of antimatter. The Gerda collaboration searches for the $0\nu\beta\beta$ decay of ^{76}Ge by operating bare germanium detectors in an active liquid argon shield. With a total exposure of $82.4 \text{ kg} \cdot \text{yr}$, we observe no signal and derive a lower half-life limit of $T_{1/2} > 0.9 \cdot 10^{26} \text{ yr}$ (90% C.L.). Our $T_{1/2}$ sensitivity assuming no signal is $1.1 \cdot 10^{26} \text{ yr}$. Combining the latter with those from other $0\nu\beta\beta$ decay searches yields a sensitivity to the effective Majorana neutrino mass of $0.07 - 0.16 \text{ eV}$.

Neutrinos were discovered in 1956 (1), but it was only at the turn of the millennium that it was experimentally proven that the three known neutrino types (flavors) ν_α ($\alpha = e, \mu, \tau$) can convert into one another (2–4). These flavor oscillations are possible only if neutrinos have nonzero mass which is currently the only established contradiction to the standard

model (SM) of particle physics. From tritium β decay experiments (5, 6) and cosmological observations (7) we know that their masses are very small - less than 10^{-5} of the electron mass. Neutrinos are the only fundamental spin-1/2 particles (fermions) without electric charge. As a consequence, they might be Majorana fermions (8) - particles identical to

their antiparticles. This is a key ingredient of some explanations for why matter is so much more abundant than antimatter in today's universe and why neutrinos are so much lighter than the other elementary particles (9).

Majorana neutrinos would lead to nuclear decays that violate lepton number conservation and are therefore forbidden in the SM of particle physics. The so-called neutrinoless double- β ($0\nu\beta\beta$) decay transforms simultaneously two neutrons inside a nucleus into two protons with an emission of two electrons, see Fig. 1. The SM-allowed double- β ($2\nu\beta\beta$) decay occurs with an emission of two electrons and two antineutrinos. In the $0\nu\beta\beta$ decay, the two electrons together carry the available decay energy ($Q_{\beta\beta}$) and the resulting mono-energetic signal is the chief experimental signature. A positive detection of this process would imply the first observation of a matter-creating process, without the balancing emission of antimatter, and would establish the Majorana nature of neutrinos (10, 11).

We report here on the search for the $0\nu\beta\beta$ decay $^{76}\text{Ge} \rightarrow ^{76}\text{Se} + 2e^-$ [$Q_{\beta\beta} = 2039.061 \pm 0.007$ keV (12)] with the GERmanium Detector Array (Gerda). Unlike previous experiments, Gerda surpasses the sensitivity for a $0\nu\beta\beta$ decay half-life of $T_{1/2} \sim 10^{26}$ yr (90% C.L.) and operates in a background-free regime such that the expected number of background events is less than 1 in the energy region of interest at the final exposure (13); here the sensitivity is defined as the median limit expected from many repetitions of the experiment assuming no signal.

This achievement together with the excellent energy resolution of Ge detectors is crucial in reaching a regime where it would be possible to detect a nonzero signal for the decay.

The Gerda experimental design was guided by the requirement to reduce interfering signals from naturally occurring radioactivity and from cosmic rays to negligible levels. The Ge detectors are made from high purity (99.9999%) Ge material that is enriched in the ^{76}Ge isotope from the natural abundance of 7.8% to more than 85%. The Ge detectors act as both the source and detector for the $0\nu\beta\beta$ decay, as illustrated in Fig. 1. In total, Gerda deploys 37 enriched detectors with two different geometries (coaxial and Broad Energy Ge (BEGe) detectors, see fig. S1) and with a total mass of 35.6 kg as bare crystals in 63 m³ of liquid argon (LAr). The LAr serves as high-purity shielding against radiation from radioactive decays, and it also provides cooling for the Ge diodes. Moreover, the LAr - due to its scintillation property - acts as a veto system to discard events originating from background radiation, which simultaneously deposit energy inside the Ge detectors and the adjacent LAr. The scintillation light is detected by 16 photomultipliers and wavelength shifting fibers connected

to silicon photomultipliers. A water tank encloses the LAr cryostat to further attenuate γ radiation and neutrons from the experimental environment. It also serves as a water Cherenkov detector to identify cosmic ray muons and their secondary shower particles that could mimic signal events. Gerda is operated deep underground at the Gran Sasso National Laboratories (LNGS) of INFN in Italy, at a depth of 3500 m water equivalent to reduce the cosmic ray muon flux by six orders of magnitude with respect to the earth's surface. Detailed descriptions of Phases I and II of the experiment can be found in (14, 15).

The signals of the Ge detectors are read out by low-radioactive charge sensitive amplifiers, digitized at 100 MHz sampling rate and stored for off-line analysis. Weekly calibrations with ^{228}Th sources are performed to monitor the energy scale and resolution, as well as to define and monitor the analysis cuts. The derived energy resolution, full width at half maximum (FWHM), at $Q_{\beta\beta}$ is 3.6 ± 0.1 keV for the coaxial and 3.0 ± 0.1 keV for the BEGe detectors, both corresponding to $\sigma/Q_{\beta\beta} < 10^{-3}$ ($\sigma = \text{FWHM} / 2.35$).

During physics data taking all Ge and LAr scintillation channels are read out if one or more Ge diodes detect a signal above a preset trigger threshold. Multiple detector hits are discarded as background events. Similarly, events are classified as background (Fig. 1) if at least one photoelectron is detected in the LAr within ~ 6 μs around the Ge detector signal - that is ~ 5 times the lifetime of the argon excimer observed in Gerda. Random coincidences lead to a loss of potential $0\nu\beta\beta$ signals of $(2.3 \pm 0.1)\%$. All events with a muon trigger preceding a Ge trigger by less than 10 μs are rejected with a signal loss of $< 0.1\%$. Background events from γ radiation lead often to multiple interactions separated in space, but within the same detector. The time structure of the recorded signal allows to reject this background as well as events occurring at the surface of a detector from α or β decays (Fig. 1, pulse shape discrimination, PSD). More than 95% of the background is rejected by the LAr veto and PSD (see Fig. 2) whereas 69% of the $0\nu\beta\beta$ decay events would be kept for the coaxial and 86% for the BEGe detectors. Compared to (16), the Phase II exposure has been more than doubled while improving both energy resolution (by 10%) and background rate (by $\sim 80\%$) in the coaxial detectors and maintaining the excellent energy resolution of the BEGe detectors throughout the run - altogether doubling the sensitivity to more than 10^{26} yr.

Since the outset, Gerda has adopted a rigorous blind analysis strategy to ensure an unbiased search for $0\nu\beta\beta$ decays. Events with a reconstructed energy of $Q_{\beta\beta} \pm 25$ keV are blinded, i.e., removed from the data stream, until the data selection is fixed. Figure 2 displays the energy spectra corresponding to 53.9 kg $\cdot$ yr Phase II exposure before and after

analysis cuts, including a PSD method for coaxial detectors that was not used in prior work (15). At low energies the spectrum after analysis cuts is dominated by $2\nu\beta\beta$ decays. The insets in the figure display separately the event distribution of the coaxial and of the BEGe detector data sets in the analysis window 1930-2190 keV. After unblinding, only 3 events in the coaxial and 4 events in the BEGe data sets remain in the analysis window (17). Gerda thus reaches an unprecedented low background rate of $5.7^{+4.1}_{-2.6} \cdot 10^{-4}$ counts/(keV · kg · yr) for the coaxial and $5.6^{+3.4}_{-2.4} \cdot 10^{-4}$ counts/(keV · kg · yr) for the BEGe detectors.

An unbinned maximum likelihood fit is carried out simultaneously to the different data sets (see table S3) including those from Gerda Phase I (18). In total, 82.4 kg · yr have been scrutinized for a $0\nu\beta\beta$ signal so far. The fit function (13) comprises flat distributions for the background, independent for each data set, and Gaussian distributions for a possible $0\nu\beta\beta$ signal: the mean is $Q_{\beta\beta}$, the resolutions are taken from calibration data individually for each set, and the normalizations are calculated from the searched for half-life $T_{1/2}$. A null signal maximizes the likelihood. Confidence intervals are evaluated both in the frequentist and Bayesian frameworks (15). The frequentist analysis is based on the profile likelihood method and systematic uncertainties are included as nuisance parameters with Gaussian pull terms. The derived limit of $T_{1/2} > 0.9 \cdot 10^{26}$ yr (90% C.L.) is compatible with the sensitivity assuming no signal of $1.1 \cdot 10^{26}$ yr; this is an improvement over previous experiments which had sensitivities less than 10^{26} yr. The weaker limit is a consequence of an event in the signal region at 2042.1 keV, 2.4 standard deviations (σ) away from $Q_{\beta\beta}$. The statistical analysis attributes it to background. Statistical analysis including Bayesian inference is detailed in (15).

Table 1 compares our results with those of other $0\nu\beta\beta$ decay searches. The $T_{1/2}$ sensitivities of other experiments are at most half of ours despite sometimes higher exposures; this is caused by Gerda's lower background and superior energy resolution (15). Several physical processes beyond the SM can produce $0\nu\beta\beta$ decay. Here we focus on the paradigm of the mixing of 3 light Majorana neutrinos. In this context, the half-life can be converted into a $0\nu\beta\beta$ decay strength that has the dimension of mass, denoted effective Majorana mass (19),

$$m_{\beta\beta} = \left| \sum_{i=1}^3 U_{ei}^2 m_i \right| \quad (1)$$

Nuclear structure details enter the decay rate, and uncertainties in the nuclear structure calculations result in a spread of $m_{\beta\beta}$ values for a given $T_{1/2}$ by typically a factor of 2-

3 (20). Some reported half-life limits L deviate by almost a factor of 2 from the associated sensitivity S , indicating significant under-fluctuation (CUORE, KamLAND-Zen), or upward fluctuation (EXO-200). To overcome this possible behavior of frequentist limits, we use the sensitivity to extract the constraints on $m_{\beta\beta}$ shown in Table 1. For Gerda the median limit is $m_{\beta\beta} < 0.1 - 0.23$ eV. Combining it with the sensitivities of the other searches (15) the bound tightens to $m_{\beta\beta} < 0.07 - 0.16$ eV (90% C.L.), very similar to the bound deduced by KamLAND-Zen from their $T_{1/2}$ limit (21).

Figure 3 shows the dependence of the effective Majorana mass $m_{\beta\beta}$ as a function of the lightest neutrino mass $m_{\text{light}} = \min(m_i)$, the cosmological observable of the sum of neutrino masses $\Sigma = \sum_i m_i$, and the effective neutrino mass

$$m_{\beta} = \sqrt{\sum_i |U_{ei}|^2 m_i^2} \quad (2)$$

i.e., the mass observable in single beta decays. The allowed parameter space is classified according to the ordering of the neutrino mass eigenstates as normal ($\Delta m_{31}^2 > 0$) or inverted ($\Delta m_{31}^2 < 0$). The overlap region is called quasi-degenerate; here the mass splittings are small compared to the absolute mass scale. Latest oscillation data prefer normal ordering at the 3σ level (22). Figure 3 shows that our extracted limits of $m_{\beta\beta}$ disfavor a large fraction of the parameter space of quasi-degenerate Majorana neutrino masses. The combined limit of $m_{\beta\beta} = 0.16$ eV corresponds to constraints on $m_{\text{light}} < 0.15 - 0.44$ eV, $\Sigma < 0.46 - 1.3$ eV and $m_{\beta} < 0.16 - 0.44$ eV. Direct measurements of m_{β} yield a limit of ~ 2.3 eV (5, 6). In the upcoming years the KATRIN tritium decay experiment will increase the sensitivity to ~ 0.2 eV (23). The sum of the neutrino masses influences the evolution and structure of the Universe. In the framework of the 6+1 parameter cosmological SM, the latest Planck data on the anisotropy of the cosmic microwave radiation along with baryonic acoustic oscillation data provide limits as low as $\Sigma < 0.12$ eV (95% C.L.) (24). Extended models relax these limits to < 0.37 eV (< 0.66 eV) for 1 (5) additional parameters (7).

Currently, there are no tensions among the 3 mass observables. A discovery of $0\nu\beta\beta$ decay close to the current experimental half-life sensitivity should have counterpart signals in tritium β decay and in cosmology, provided that the paradigm of 3 light Majorana neutrinos holds. In case of discrepancies with the other mass observables, a $0\nu\beta\beta$ signal would point to other lepton number violating processes. Within the framework of 3 light Majorana neutrinos, the cosmological SM and in the absence of a $0\nu\beta\beta$ decay at or close to the current sensitivity, the KATRIN experiment

would not observe a signal. Conversely, a positive measurement of $m_\beta > 0.44$ eV in KATRIN would point to Dirac neutrinos or to an incomplete understanding of the nuclear physics (20) of $0\nu\beta\beta$ decay. It also would require extensions to the current minimal cosmological model. Instead, if the cosmological limit on Σ holds, $0\nu\beta\beta$ decay experiments would have to probe a mass range $m_{\beta\beta} < 0.05$ eV which requires a half-life sensitivity of 10^{27} yr and above for a ^{76}Ge -based experiment.

The leading performance of Gerda in terms of background suppression, energy resolution and sensitivity opens the way to Legend, a next generation Ge experiment with sensitivity to half-lives of 10^{27} yr and beyond. A first phase 200 kg ^{76}Ge experiment, Legend-200 (25), is under preparation at LNGS.

REFERENCES AND NOTES

1. C. L. Cowan Jr., F. Reines, F. B. Harrison, H. W. Kruse, A. D. McGuire, Detection of the Free Neutrino: A Confirmation. *Science* **124**, 103–104 (1956).
[doi:10.1126/science.124.3212.103](https://doi.org/10.1126/science.124.3212.103) [Medline](#)
2. Y. Fukuda, T. Hayakawa, E. Ichihara, K. Inoue, K. Ishihara, H. Ishino, Y. Itow, T. Kajita, J. Kameda, S. Kasuga, K. Kobayashi, Y. Kobayashi, Y. Koshio, M. Miura, M. Nakahata, S. Nakayama, A. Okada, K. Okumura, N. Sakurai, M. Shiozawa, Y. Suzuki, Y. Takeuchi, Y. Totsuka, S. Yamada, M. Earl, A. Habig, E. Kearns, M. D. Messier, K. Scholberg, J. L. Stone, L. R. Sulak, C. W. Walter, M. Goldhaber, T. Barszczak, D. Casper, W. Gajewski, P. G. Halverson, J. Hsu, W. R. Kropp, L. R. Price, F. Reines, M. Smy, H. W. Sobel, M. R. Vagins, K. S. Ganezer, W. E. Keig, R. W. Ellsworth, S. Tasaka, J. W. Flanagan, A. Kibayashi, J. G. Learned, S. Matsuno, V. J. Stenger, D. Takemori, T. Ishii, J. Kanzaki, T. Kobayashi, S. Mine, K. Nakamura, K. Nishikawa, Y. Oyama, A. Sakai, M. Sakuda, O. Sasaki, S. Echigo, M. Kohama, A. T. Suzuki, T. J. Haines, E. Blaufuss, B. K. Kim, R. Sanford, R. Svoboda, M. L. Chen, Z. Conner, J. A. Goodman, G. W. Sullivan, J. Hill, C. K. Jung, K. Martens, C. Mauger, C. McGrew, E. Sharkey, B. Viren, C. Yanagisawa, W. Doki, K. Miyano, H. Okazawa, C. Saji, M. Takahata, Y. Nagashima, M. Takita, T. Yamaguchi, M. Yoshida, S. B. Kim, M. Etoh, K. Fujita, A. Hasegawa, T. Hasegawa, S. Hatakeyama, T. Iwamoto, M. Koga, T. Maruyama, H. Ogawa, J. Shirai, A. Suzuki, F. Tushima, M. Koshihara, M. Nemoto, K. Nishijima, T. Futagami, Y. Hayato, Y. Kanaya, K. Kaneyuki, Y. Watanabe, D. Kielczewska, R. A. Doyle, J. S. George, A. L. Stachyra, L. L. Wai, R. J. Wilkes, K. K. Young, Evidence for Oscillation of Atmospheric Neutrinos. *Phys. Rev. Lett.* **81**, 1562–1567 (1998).
[doi:10.1103/PhysRevLett.81.1562](https://doi.org/10.1103/PhysRevLett.81.1562)
3. Q. R. Ahmad, R. C. Allen, T. C. Andersen, J. D. Anglin, J. C. Barton, E. W. Beier, M. Bercovitch, J. Bigu, S. D. Biller, R. A. Black, I. Blevins, R. J. Boardman, J. Boger, E. Bonvin, M. G. Boulay, M. G. Bowler, T. J. Bowles, S. J. Brice, M. C. Browne, T. V. Bullard, G. Bühler, J. Cameron, Y. D. Chan, H. H. Chen, M. Chen, X. Chen, B. T. Cleveland, E. T. Clifford, J. H. Cowan, D. F. Cowen, G. A. Cox, X. Dai, F. Dalnoki-Veress, W. F. Davidson, P. J. Doe, G. Doucas, M. R. Dragowsky, C. A. Duba, F. A. Duncan, M. Dunford, J. A. Dunmore, E. D. Earle, S. R. Elliott, H. C. Evans, G. T. Ewan, J. Farine, H. Fergani, A. P. Ferraris, R. J. Ford, J. A. Formaggio, M. M. Fowler, K. Frame, E. D. Frank, W. Frati, N. Gagnon, J. V. Germani, S. Gil, K. Graham, D. R. Grant, R. L. Hahn, A. L. Hallin, E. D. Hallman, A. S. Hamer, A. A. Hamian, W. B. Handler, R. U. Haq, C. K. Hargrove, P. J. Harvey, R. Hazama, K. M. Heeger, W. J. Heintzelman, J. Heise, R. L. Helmer, J. D. Hepburn, H. Heron, J. Hewett, A. Hime, M. Howe, J. G. Hykawy, M. C. Isaac, P. Jagam, N. A. Jelley, C. Jillings, G. Jonkmans, K. Kazkaz, P. T. Keener, J. R. Klein, A. B. Knox, R. J. Komar, R. Kouzes, T. Kutter, C. C. Kyba, J. Law, I. T. Lawson, M. Lay, H. W. Lee, K. T. Lesko, J. R. Leslie, I. Levine, W. Locke, S. Luoma, J. Lyon, S. Majerus, H. B. Mak, J. Maneira, J. Manor, A. D. Marino, N. McCauley, A. B. McDonald, D. S. McDonald, K. McFarlane, G. McGregor, R. Meijer Drees, C. Mifflin, G. G. Miller, G. Milton, B. A. Moffat, M. Moorhead, C. W. Nally, M. S. Neubauer, F. M. Newcomer, H. S. Ng, A. J. Noble, E. B. Norman, V. M. Novikov, M. O'Neill, C. E. Okada, R. W. Ollerhead, M. Omori, J. L. Orrell, S. M. Oser, A. W. Poon, T. J. Radcliffe, A. Roberge, B. C. Robertson, R. G. Robertson, S. S. Rosendahl, J. K. Rowley, V. L. Rusu, E. Saettler, K. K. Schaffer, M. H. Schwendener, A. Schülke, H. Seifert, M. Shatkay, J. J. Simpson, C. J. Sims, D. Sinclair, P. Skensved, A. R. Smith, M. W. Smith, T. Spreitzer, N. Starinsky, T. D. Steiger, R. G. Stokstad, L. C. Stonehill, R. S. Storey, B. Sur, R. Tafirout, N. Tagg, N. W. Tanner, R. K. Taplin, M. Thorman, P. M. Thornewell, P. T. Trent, Y. I. Tserkovnyak, R. Van Berg, R. G. Van de Water, C. J. Virtue, C. E. Waltham, J. X. Wang, D. L. Wark, N. West, J. B. Wilhelmy, J. F. Wilkerson, J. R. Wilson, P. Wittich, J. M. Wouters, M. Yeh, Direct evidence for neutrino flavor transformation from neutral-current interactions in the Sudbury Neutrino Observatory. *Phys. Rev. Lett.* **89**, 011301 (2002).
[doi:10.1103/PhysRevLett.89.011301](https://doi.org/10.1103/PhysRevLett.89.011301) [Medline](#)
4. K. Eguchi, S. Enomoto, K. Furuno, J. Goldman, H. Hanada, H. Ikeda, K. Ikeda, K. Inoue, K. Ishihara, W. Itoh, T. Iwamoto, T. Kawaguchi, T. Kawashima, H. Kinoshita, Y. Kishimoto, M. Koga, Y. Koseki, T. Maeda, T. Mitsui, M. Motoki, K. Nakajima, M. Nakajima, T. Nakajima, H. Ogawa, K. Owada, T. Sakabe, I. Shimizu, J. Shirai, F. Suekane, A. Suzuki, K. Tada, O. Tajima, T. Takayama, K. Tamae, H. Watanabe, J. Busenitz, Z. Djuric, K. McKinny, D.-M. Mei, A. Piepke, E. Yakushev, B. E. Berger, Y. D. Chan, M. P. Decowski, D. A. Dwyer, S. J. Freedman, Y. Fu, B. K. Fujikawa, K. M. Heeger, K. T. Lesko, K.-B. Luk, H. Murayama, D. R. Nygren, C. E. Okada, A. W. P. Poon, H. M. Steiner, L. A. Winslow, G. A. Horton-Smith, R. D. McKeown, J. Ritter, B. Tipton, P. Vogel, C. E. Lane, T. Miletic, P. W. Gorham, G. Guillian, J. G. Learned, J. Maricic, S. Matsuno, S. Pakvasa, S. Dazeley, S. Hatakeyama, M. Murakami, R. C. Svoboda, B. D. Dieterle, M. DiMauro, J. Detwiler, G. Gratta, K. Ishii, N. Tolich, Y. Uchida, M. Batygov, W. Bugg, H. Cohn, Y. Efremenko, Y. Kamyshev, A. Kozlov, Y. Nakamura, L. De Braeckeleer, C. R. Gould, H. J. Karwowski, D. M. Markoff, J. A. Messimore, K. Nakamura, R. M. Rohm, W. Tornow, A. R. Young, Y.-F. Wang, First results from KamLAND: Evidence for reactor antineutrino disappearance. *Phys. Rev. Lett.* **90**, 021802 (2003).
[doi:10.1103/PhysRevLett.90.021802](https://doi.org/10.1103/PhysRevLett.90.021802) [Medline](#)
5. C. Kraus, B. Bornschein, L. Bornschein, J. Bonn, B. Flatt, A. Kovalik, B. Ostrick, E. W. Otten, J. P. Schall, T. Thümmel, C. Weinheimer, Final results from phase II of the Mainz neutrino mass search in tritium β decay. *Eur. Phys. J. C* **40**, 447–468 (2005).
[doi:10.1140/epjc/s2005-02139-7](https://doi.org/10.1140/epjc/s2005-02139-7)
6. V. N. Aseev, A. I. Belev, A. I. Berlev, E. V. Geraskin, A. A. Golubev, N. A. Likhovid, V. M. Lobashev, A. A. Nozik, V. S. Pantuev, V. I. Parfenov, A. K. Skasyrskaia, F. V. Tkachov, S. V. Zadorozhny, Upper limit on the electron antineutrino mass from the Troitsk experiment. *Phys. Rev. D* **84**, 112003 (2011).
[doi:10.1103/PhysRevD.84.112003](https://doi.org/10.1103/PhysRevD.84.112003)
7. M. Tanabashi, K. Hagiwara, K. Hikasa, K. Nakamura, Y. Sumino, F. Takahashi, J. Tanaka, K. Agashe, G. Aielli, C. Amsler, M. Antonelli, D. M. Asner, H. Baer, S. Banerjee, R. M. Barnett, T. Basaglia, C. W. Bauer, J. J. Beatty, V. I. Belousov, J. Beringer, S. Bethke, A. Bettini, H. Bichsel, O. Biebel, K. M. Black, E. Blucher, O. Buchmüller, V. Burkert, M. A. Bychkov, R. N. Cahn, M. Carena, A. Ceccucci, A. Cerri, D. Chakraborty, M.-C. Chen, R. S. Chivukula, G. Cowan, O. Dahl, G. D'Ambrosio, T. Damour, D. de Florian, A. de Gouvêa, T. DeGrand, P. de Jong, G. Dissertori, B. A. Dobrescu, M. D'Onofrio, M. Doser, M. Drees, H. K. Dreiner, D. A. Dwyer, P. Eerola, S. Eidelman, J. Ellis, J. Erler, V. V. Ezhela, W. Fetscher, B. D. Fields, R. Firestone, B. Foster, A. Freitas, H. Gallagher, L. Garren, H.-J. Gerber, G. Gerbier, T. Gershon, Y. Gershtein, T. Gherghetta, A. A. Godizov, M. Goodman, C. Grab, A. V. Gritsan, C. Grojean, D. E. Groom, M. Grünewald, A. Gurtu, T. Gutsche, H. E. Haber, C. Hanhart, S. Hashimoto, Y. Hayato, K. G. Hayes, A. Hebecker, S. Heinemeyer, B. Heltsley, J. J. Hernández-Rey, J. Hisano, A. Höcker, J. Holder, A. Holtkamp, T. Hyodo, K. D. Irwin, K. F. Johnson, M. Kado, M. Karliner, U. F. Katz, S. R. Klein, E. Klempf, R. V. Kowalewski, F. Krauss, M. Kreps, B. Krusche, Y. V. Kuyanov, Y. Kwon, O. Lahav, J. Laiho, J. Lesgourgues, A. Liddle, Z. Ligeti, C.-J. Lin, C. Lippmann, T. M. Liss, L. Littenberg, K. S. Lugovsky, S. B. Lugovsky, A. Lusiani, Y. Makida, F. Maltoni, T. Mannel, A. V. Manohar, W. J. Marciano, A. D. Martin, A. Masoni, J. Matthews, U.-G. Meißner, D. Milstead, R. E. Mitchell, K. Mönig, P. Molaro, F. Moortgat, M. Moskvic, H. Murayama, M. Narain, P. Nason, S. Navas, M. Neubert, P. Nevski, Y. Nir, K. A. Olive, S. Pagan Griso, J. Parsons, C. Patrignani, J. A. Peacock, M. Pennington, S. T. Petcov, V. A. Petrov, E. Pianori, A.

- Piepeke, A. Pomarol, A. Quad, J. Rademacker, G. Raffelt, B. N. Ratcliff, P. Richardson, A. Ringwald, S. Roesler, S. Rolli, A. Romaniouk, L. J. Rosenberg, J. L. Rosner, G. Rybka, R. A. Rytin, C. T. Sachrajda, Y. Sakai, G. P. Salam, S. Sarkar, F. Sauli, O. Schneider, K. Scholberg, A. J. Schwartz, D. Scott, V. Sharma, S. R. Sharpe, T. Shutt, M. Silari, T. Sjöstrand, P. Skands, T. Skwarnicki, J. G. Smith, G. F. Smoot, S. Spanier, H. Spieler, C. Spiering, A. Stahl, S. L. Stone, T. Sumiyoshi, M. J. Syphers, K. Terashi, J. Terning, U. Thoma, R. S. Thorne, L. Tiator, M. Titov, N. P. Tkachenko, N. A. Törnqvist, D. R. Tovey, G. Valencia, R. Van de Water, N. Varelas, G. Venanzoni, L. Verde, M. G. Vincter, P. Vogel, A. Vogt, S. P. Wakely, W. Walkowiak, C. W. Walter, D. Wands, D. R. Ward, M. O. Wascko, G. Weiglein, D. H. Weinberg, E. J. Weinberg, M. White, L. R. Wiencke, S. Willocq, C. G. Wohl, J. Womersley, C. L. Woody, R. L. Workman, W.-M. Yao, G. P. Zeller, O. V. Zenin, R.-Y. Zhu, S.-L. Zhu, F. Zimmermann, P. A. Zyla, J. Anderson, L. Fuller, V. S. Lugovsky, P. Schaffner, Review of Particle Physics. *Phys. Rev. D* **98**, 030001 (2018). doi:10.1103/PhysRevD.98.030001
8. E. Majorana, Teoria simmetrica dell'elettrone e del positrone. *Nuovo Cim.* **14**, 171–184 (1937). doi:10.1007/BF02961314
9. R. N. Mohapatra, A. Y. Smirnov, Neutrino Mass and New Physics. *Annu. Rev. Nucl. Part. Sci.* **56**, 569–628 (2006). doi:10.1146/annurev.nucl.56.080805.140534
10. J. Schechter, J. W. F. Valle, Neutrino masses in SU(2) $\otimes$ U(1) theories. *Phys. Rev. D* **22**, 2227–2235 (1980). doi:10.1103/PhysRevD.22.2227
11. S. Dell'Oro, S. Marcocci, F. Vissani, Testing creation of matter with neutrinoless double beta decay. *PoS(NEUTEL2017) 030* (2018); <https://pos.sissa.it/307/030/pdf>.
12. B. J. Mount, M. Redshaw, E. G. Myers, Double- β -decay Q values of ^{74}Se and ^{76}Ge . *Phys. Rev. C* **81**, 032501 (2010). doi:10.1103/PhysRevC.81.032501
13. GERDA Collaboration, Background-free search for neutrinoless double- β decay of ^{76}Ge with GERDA. *Nature* **544**, 47–52 (2017). doi:10.1038/nature21717 Medline
14. M. Agostini, A. M. Bakalyarov, M. Balata, I. Barabanov, L. Baudis, C. Bauer, E. Bellotti, S. Belogurov, S. T. Belyaev, G. Benato, A. Bettini, L. Bezrukov, T. Bode, D. Borowicz, V. Brudanin, R. Brugnera, A. Caldwell, C. Cattadori, A. Chernogorov, V. D'Andrea, E. V. Demidova, N. Di Marco, A. Domula, E. Doroshkevich, V. Egorov, R. Falkenstein, N. Frodyma, A. Gangapshv, A. Garfagnini, P. Grabmayr, V. Gurentsov, K. Gusev, J. Hakenmüller, A. Hegai, M. Heisel, S. Hemmer, R. Hiller, W. Hofmann, M. Hult, L. V. Inzhechik, L. Ioannucci, J. Janicskó Csáthy, J. Jochum, M. Junker, V. Kazalov, Y. Kermaidic, T. Kihm, I. V. Kirpichnikov, A. Kirsch, A. Kish, A. Klimenko, R. Kneiβl, K. T. Knöpfle, O. Kochetov, V. N. Kornoukhov, V. V. Kuzminov, M. Laubenstein, A. Lazzaro, V. I. Lebedev, B. Lehnert, M. Lindner, I. Lippi, A. Lubashevskiy, B. Lubsandorzhiev, G. Lutter, C. Macolino, B. Majorovits, W. Maneschg, E. Medinaceli, M. Miloradovic, R. Mingazheva, M. Misiaszek, P. Moseev, I. Nemchenok, S. Nisi, K. Panas, L. Pandola, K. Pelczar, A. Pullia, C. Ransom, S. Riboldi, N. Rumyantseva, C. Sada, F. Salamida, M. Salathe, C. Schmitt, B. Schneider, S. Schönert, J. Schreiner, A.-K. Schütz, O. Schulz, B. Schwingenheuer, O. Selivanenko, E. Shevchik, M. Shirchenko, H. Simgen, A. Smolnikov, L. Stanco, L. Vanhoefer, A. A. Vasenko, A. Veresnikova, K. von Sturm, V. Wagner, A. Wegmann, T. Wester, C. Wiesinger, M. Wojcik, E. Yanovich, I. Zhitnikov, S. V. Zhukov, D. Zinatulina, A. J. Zsigmond, K. Zuber, G. Zuzel, Upgrade for Phase II of the Gerda experiment. *Eur. Phys. J. C* **78**, 388 (2018). doi:10.1140/epjc/s10052-018-5812-2
15. See supplementary materials.
16. M. Agostini, A. M. Bakalyarov, M. Balata, I. Barabanov, L. Baudis, C. Bauer, E. Bellotti, S. Belogurov, A. Bettini, L. Bezrukov, J. Biernat, T. Bode, D. Borowicz, V. Brudanin, R. Brugnera, A. Caldwell, C. Cattadori, A. Chernogorov, T. Comellato, V. D'Andrea, E. V. Demidova, N. Di Marco, A. Domula, E. Doroshkevich, V. Egorov, R. Falkenstein, A. Gangapshv, A. Garfagnini, P. Grabmayr, V. Gurentsov, K. Gusev, J. Hakenmüller, A. Hegai, M. Heisel, S. Hemmer, R. Hiller, W. Hofmann, M. Hult, L. V. Inzhechik, J. Janicskó Csáthy, J. Jochum, M. Junker, V. Kazalov, Y. Kermaidic, T. Kihm, I. V. Kirpichnikov, A. Kirsch, A. Kish, A. Klimenko, R. Kneiβl, K. T. Knöpfle, O. Kochetov, V. N. Kornoukhov, V. V. Kuzminov, M. Laubenstein, A. Lazzaro, M. Lindner, I. Lippi, A. Lubashevskiy, B. Lubsandorzhiev, G. Lutter, C. Macolino, B. Majorovits, W. Maneschg, M. Miloradovic, R. Mingazheva, M. Misiaszek, P. Moseev, I. Nemchenok, K. Panas, L. Pandola, K. Pelczar, L. Pertoldi, A. Pullia, C. Ransom, S. Riboldi, N. Rumyantseva, C. Sada, F. Salamida, C. Schmitt, B. Schneider, S. Schönert, A.-K. Schütz, O. Schulz, B. Schwingenheuer, O. Selivanenko, E. Shevchik, M. Shirchenko, H. Simgen, A. Smolnikov, L. Stanco, L. Vanhoefer, A. A. Vasenko, A. Veresnikova, K. von Sturm, V. Wagner, A. Wegmann, T. Wester, C. Wiesinger, M. Wojcik, E. Yanovich, I. Zhitnikov, S. V. Zhukov, D. Zinatulina, A. J. Zsigmond, K. Zuber, G. Zuzel, Improved Limit on Neutrinoless Double- β Decay of ^{76}Ge from GERDA Phase II. *Phys. Rev. Lett.* **120**, 132503 (2018). doi:10.1103/PhysRevLett.120.132503 Medline
17. Three ± 5 keV wide intervals at 2104 keV and 2119 keV, the position of known γ lines, and $Q_{\beta\beta}$ are excluded for the calculation of the background rate. The limit calculation, however, includes the interval $Q_{\beta\beta} \pm 5$ keV.
18. M. Agostini, M. Allardt, E. Andreotti, A. M. Bakalyarov, M. Balata, I. Barabanov, M. Barnabé Heider, N. Barros, L. Baudis, C. Bauer, N. Becerici-Schmidt, E. Bellotti, S. Belogurov, S. T. Belyaev, G. Benato, A. Bettini, L. Bezrukov, T. Bode, V. Brudanin, R. Brugnera, D. Budjáš, A. Caldwell, C. Cattadori, A. Chernogorov, F. Cossavella, E. V. Demidova, A. Domula, V. Egorov, R. Falkenstein, A. Ferella, K. Freund, N. Frodyma, A. Gangapshv, A. Garfagnini, C. Gotti, P. Grabmayr, V. Gurentsov, K. Gusev, K. K. Guthikonda, W. Hampel, A. Hegai, M. Heisel, S. Hemmer, G. Heusser, W. Hofmann, M. Hult, L. V. Inzhechik, L. Ioannucci, J. Janicskó Csáthy, J. Jochum, M. Junker, T. Kihm, I. V. Kirpichnikov, A. Kirsch, A. Klimenko, K. T. Knöpfle, O. Kochetov, V. N. Kornoukhov, V. V. Kuzminov, M. Laubenstein, A. Lazzaro, V. I. Lebedev, B. Lehnert, H. Y. Liao, M. Lindner, I. Lippi, X. Liu, A. Lubashevskiy, B. Lubsandorzhiev, G. Lutter, C. Macolino, A. A. Machado, B. Majorovits, W. Maneschg, M. Misiaszek, I. Nemchenok, S. Nisi, C. O'Shaughnessy, L. Pandola, K. Pelczar, G. Pessina, A. Pullia, S. Riboldi, N. Rumyantseva, C. Sada, M. Salathe, C. Schmitt, J. Schreiner, O. Schulz, B. Schwingenheuer, S. Schönert, E. Shevchik, M. Shirchenko, H. Simgen, A. Smolnikov, L. Stanco, H. Strecker, M. Tarka, C. A. Ur, A. A. Vasenko, O. Volynets, K. von Sturm, V. Wagner, M. Walter, A. Wegmann, T. Wester, M. Wojcik, E. Yanovich, P. Zavarise, I. Zhitnikov, S. V. Zhukov, D. Zinatulina, K. Zuber, G. Zuzel, Results on neutrinoless double- β decay of ^{76}Ge from phase I of the GERDA experiment. *Phys. Rev. Lett.* **111**, 122503 (2013). doi:10.1103/PhysRevLett.111.122503 Medline
19. The unitary 3×3 matrix $U_{\alpha i}$ relates neutrino flavor states ν_{α} ($\alpha = e, \mu, \tau$) and mass eigenstates ν_i ($i = 1, 2, 3$). The absolute neutrino masses are still unknown, but two squared neutrino mass differences Δm_{21}^2 and $|\Delta m_{31}^2|$, ($\Delta m_{ij}^2 = m_i^2 - m_j^2$), are known with increasing precision from neutrino oscillation experiments (22).
20. J. Engel, J. Menéndez, Status and future of nuclear matrix elements for neutrinoless double-beta decay: A review. *Rep. Prog. Phys.* **80**, 046301 (2017). doi:10.1088/1361-6633/aa5bc5 Medline
21. A. Gando, Y. Gando, T. Hachiya, A. Hayashi, S. Hayashida, H. Ikeda, K. Inoue, K. Ishidoshio, Y. Karino, M. Koga, S. Matsuda, T. Mitsui, K. Nakamura, S. Obara, T. Oura, H. Ozaki, I. Shimizu, Y. Shirahata, J. Shirai, A. Suzuki, T. Takai, K. Tamae, Y. Teraoka, K. Ueshima, H. Watanabe, A. Kozlov, Y. Takemoto, S. Yoshida, K. Fushimi, T. I. Banks, B. E. Berger, B. K. Fujikawa, T. O'Donnell, L. A. Winslow, Y. Efremenko, H. J. Karwowski, D. M. Markoff, W. Tornow, J. A. Detwiler, S. Enomoto, M. P. Decowski, Search for Majorana Neutrinos Near the Inverted Mass Hierarchy Region with KamLAND-Zen. *Phys. Rev. Lett.* **117**, 082503 (2016). doi:10.1103/PhysRevLett.117.082503 Medline
22. I. Esteban, M. C. Gonzalez-Garcia, A. Hernandez-Cabezudo, M. Maltoni, T. Schwetz, Global analysis of three-flavour neutrino oscillations: Synergies and tensions in the determination of θ_{23} , δ_{CP} , and the mass ordering. *J. High Energy Phys.* **2019**, 106 (2019). doi:10.1007/JHEP01(2019)106
23. J. Angrik et al., KATRIN Design Report 2004 (2005); www.katrin.kit.edu/publikationen/DesignReport2004-12Jan2005.pdf.
24. Planck Collaboration, Planck 2018 results. VI. Cosmological parameters. [arXiv:1807.06209](https://arxiv.org/abs/1807.06209) [astro-ph.CO] (17 July 2018).
25. N. Abgrall, A. Abramov, N. Abrosimov, I. Abt, M. Agostini, M. Agartioglu, A. Ajjaq, S. I. Alvis, F. T. Avignone III, X. Bai, M. Balata, I. Barabanov, A. S. Barabash, P. J.

- Barton, L. Baudis, L. Bezrukov, T. Bode, A. Bolozdynya, D. Borowicz, A. Boston, H. Boston, S. T. P. Boyd, R. Breier, V. Brudanin, R. Brugnera, M. Busch, M. Buuck, A. Caldwell, T. S. Caldwell, T. Camellato, M. Carpenter, C. Cattadori, J. Cederkäll, Y.-D. Chan, S. Chen, A. Chernogorov, C. D. Christofferson, P.-H. Chu, R. J. Cooper, C. Cuesta, E. V. Demidova, Z. Deng, M. Deniz, J. A. Detwiler, N. Di Marco, A. Domula, Q. Du, Y. Efremenko, V. Egorov, S. R. Elliott, D. Fields, F. Fischer, A. Galindo-Uribarri, A. Gangapshev, A. Garfagnini, T. Gilliss, M. Giordano, G. K. Giovanetti, M. Gold, P. Golubev, C. Gooch, P. Grabmayr, M. P. Green, J. Gruszko, I. S. Guinn, V. E. Guiseppe, V. Gurentsov, Y. Gurov, K. Gusev, J. Hakenmüller, L. Harkness-Brennan, Z. R. Harvey, C. R. Haufe, L. Hauertmann, D. Heglund, L. Hehn, A. Heinz, R. Hiller, J. Hinton, R. Hodak, W. Hofmann, S. Howard, M. A. Howe, M. Hult, L. V. Inzhchik, J. J. Csáthy, R. Janssens, M. Ješkovský, J. Jochum, H. T. Johansson, D. Judson, M. Junker, J. Kaizer, K. Kang, V. Kazalov, Y. Kermadic, F. Kiessling, A. Kirsch, A. Kish, A. Klimenko, K. T. Knöpfle, O. Kochetov, S. I. Konovalov, I. Kontul, V. N. Kornoukhov, T. Kraetzschmar, K. Kröninger, A. Kumar, V. V. Kuzminov, K. Lang, M. Laubenstein, A. Lazzaro, Y. L. Li, Y.-Y. Li, H. B. Li, S. T. Lin, M. Lindner, I. Lippi, S. K. Liu, X. Liu, J. Liu, D. Loomba, A. Lubashevskiy, B. Lubsandorzhiev, G. Lutter, H. Ma, B. Majorovits, F. Mamedov, R. D. Martin, R. Massarczyk, J. A. J. Matthews, N. McFadden, D.-M. Mei, H. Mei, S. J. Meijer, D. Mengoni, S. Mertens, W. Miller, M. Miloradovic, R. Mingazheva, M. Misiaszek, P. Moseev, J. Myslik, I. Nemchenok, T. Nilsson, P. Nolan, C. O'Shaughnessy, G. Othman, K. Panas, L. Pandola, L. Papp, K. Pelczar, D. Peterson, W. Pettus, A. W. P. Poon, P. P. Povinec, A. Pullia, X. C. Quintana, D. C. Radford, J. Rager, C. Ransom, F. Recchia, A. L. Reine, S. Riboldi, K. Rielage, S. Rozov, N. W. Rouf, E. Rukhadze, N. Rumyantseva, R. Saakyan, E. Sala, F. Salamida, V. Sandukovsky, G. Savard, S. Schönert, A.-K. Schütz, O. Schulz, M. Schuster, B. Schwingenheuer, O. Selivanenko, B. Sevdá, B. Shanks, E. Shevchik, M. Shirchenko, F. Simkovic, L. Singh, V. Singh, M. Skorokhvatov, K. Smolek, A. Smolnikov, A. Sonay, M. Spavorova, I. Stekl, D. Stukov, D. Tedeschi, J. Thompson, T. Van Wechel, R. L. Varner, A. A. Vasenko, S. Vasilyev, A. Veresnikova, K. Vetter, K. von Sturm, K. Vorren, M. Wagner, G.-J. Wang, D. Waters, W.-Z. Wei, T. Wester, B. R. White, C. Wiesinger, J. F. Wilkerson, M. Willers, C. Wiseman, M. Wojcik, H. T. Wong, J. Wyenberg, W. Xu, E. Yakushev, G. Yang, C.-H. Yu, Q. Yue, V. Yumatov, J. Zeman, Z. Zeng, I. Zhitnikov, B. Zhu, D. Zinatulina, A. Zschocke, A. J. Zsigmond, K. Zuber, G. Zuzel, *AIP Conf. Proc.* **1894**, 020027 (2017). [doi:10.1063/1.5007652](https://doi.org/10.1063/1.5007652)
26. GERDA Collaboration, Dataset for “Probing Majorana neutrinos with double- β decay” (2019); Zenodo, <http://doi.org/10.5281/zenodo.339351>.
27. S. I. Alvis *et al.*, A Search for Neutrinoless Double- β Decay in ^{76}Ge with 26 kg-yr of Exposure from the MAJORANA DEMONSTRATOR. *Phys. Rev. C* **100**, 025501 (2019). [doi:10.1103/PhysRevC.100.025501](https://doi.org/10.1103/PhysRevC.100.025501)
28. O. Azzolini, M. T. Barrera, J. W. Beeman, F. Bellini, M. Beretta, M. Biassoni, C. Brofferio, C. Bucci, L. Canonica, S. Capelli, L. Cardani, P. Carniti, N. Casali, L. Cassina, M. Clemenza, O. Cremonesi, A. Cruciani, A. D'Addabbo, I. Dafinei, S. Di Domizio, F. Ferroni, L. Gironi, A. Giuliani, P. Gorla, C. Gotti, G. Keppel, L. Marini, M. Martinez, S. Morganti, S. Nagorny, M. Nastasi, S. Nisi, C. Nones, D. Orlandi, L. Pagnanini, M. Pallavicini, V. Palmieri, L. Pattavina, M. Pavan, G. Pessina, V. Pettinacci, S. Pirro, S. Pozzi, E. Previtali, A. Puiu, F. Reindl, C. Rusconi, K. Schäffner, C. Tomei, M. Vignati, A. S. Zolotarova, First Result on the Neutrinoless Double- β Decay of ^{82}Se with CUPID-0. *Phys. Rev. Lett.* **120**, 232502 (2018). [doi:10.1103/PhysRevLett.120.232502](https://doi.org/10.1103/PhysRevLett.120.232502) [Medline](https://arxiv.org/abs/1805.02550)
29. C. Alduino, F. Alessandria, K. Alfonso, E. Andreotti, C. Arnaboldi, F. T. Avignone, O. Azzolini, M. Balata, I. Bandac, T. I. Banks, G. Bari, M. Barucci, J. W. Beeman, F. Bellini, G. Benato, A. Bersani, D. Biare, M. Biassoni, F. Bragazzi, A. Branca, C. Brofferio, A. Bryant, A. Buccheri, C. Bucci, C. Bulfon, A. Camacho, A. Caminata, L. Canonica, X. G. Cao, S. Capelli, M. Capodiferro, L. Cappelli, L. Cardani, M. Cariello, P. Carniti, M. Carrettoni, N. Casali, L. Cassina, R. Cereseto, G. Ceruti, A. Chiarini, D. Chiesa, N. Chott, M. Clemenza, D. Conventi, S. Copello, C. Cosmelli, O. Cremonesi, C. Crescentini, R. J. Creswick, J. S. Cushman, A. D'Addabbo, D. D'Aguzzo, I. Dafinei, V. Datskov, C. J. Davis, F. Del Corso, S. Dell'Oro, M. M. Deninno, S. Di Domizio, M. L. Di Vacri, L. Di Paolo, A. Drobizhev, L. Ejzak, R. Faccini, D. Q. Fang, M. Faverzani, E. Ferri, F. Ferroni, E. Fiorini, M. A. Franceschi, S. J. Freedman, B. K. Fujikawa, R. Gaigher, A. Giachero, L. Gironi, A. Giuliani, L. Gladstone, J. Goett, P. Gorla, C. Gotti, C. Guandalini, M. Guerzoni, T. D. Gutierrez, E. E. Haller, K. Han, E. V. Hansen, K. M. Heeger, R. Hennings-Yeomans, K. P. Hickerson, H. Z. Huang, M. Iannone, L. Ioannucci, R. Kadel, G. Keppel, L. Kogler, Y. G. Kolomensky, A. Leder, C. Ligi, K. E. Lim, X. Liu, Y. G. Ma, C. Maiano, M. Maino, L. Marini, M. Martinez, C. Martinez Amaya, R. H. Maruyama, Y. Mei, N. Moggi, S. Morganti, P. J. Mosteiro, S. S. Nagorny, T. Napolitano, M. Nastasi, S. Nisi, C. Nones, E. B. Norman, V. Novati, A. Nucciotti, I. Nutini, T. O'Donnell, M. Olcese, E. Olivieri, F. Orio, D. Orlandi, J. L. Ouellet, C. E. Pagliarone, M. Pallavicini, V. Palmieri, L. Pattavina, M. Pavan, M. Pedretti, R. Pedrotta, A. Pelosi, G. Pessina, V. Pettinacci, G. Piperno, C. Pira, S. Pirro, S. Pozzi, E. Previtali, F. Reindl, F. Rimondi, L. Riseigari, C. Rosenfeld, C. Rossi, C. Rusconi, M. Sakai, E. Sala, C. Salvioni, S. Sangiorgio, D. Santone, D. Schaeffer, B. Schmidt, J. Schmidt, N. D. Scielzo, V. Singh, M. Sisti, A. R. Smith, F. Stivanello, L. Taffarello, L. Tatananni, M. Tenconi, F. Terranova, M. Tessaro, C. Tomei, G. Ventura, M. Vignati, S. L. Wagaarachchi, J. Wallig, B. S. Wang, H. W. Wang, B. Welliver, J. Wilson, K. Wilson, L. A. Winslow, T. Wise, L. Zanotti, C. Zarra, G. Q. Zhang, B. X. Zhu, S. Zimmermann, S. Zucchelli, First Results from CUORE: A Search for Lepton Number Violation via $0\nu\beta\beta$ Decay of ^{130}Te . *Phys. Rev. Lett.* **120**, 132501 (2018). [doi:10.1103/PhysRevLett.120.132501](https://doi.org/10.1103/PhysRevLett.120.132501) [Medline](https://arxiv.org/abs/1805.02550)
30. J. B. Albert, G. Anton, I. Badhrees, P. S. Barbeau, R. Bayerlein, D. Beck, V. Belov, M. Breidenbach, T. Brunner, G. F. Cao, W. R. Cen, C. Chambers, B. Cleveland, M. Coon, A. Craycraft, W. Cree, T. Daniels, M. Danilov, S. J. Daugherty, J. Daughettee, J. Davis, S. Delaquis, A. Der Mesrobian-Kabakian, R. DeVoe, T. Didberidze, J. Dilling, A. Dolgolenko, M. J. Dolinski, W. Fairbank, J. Farine, S. Feyzbakhsh, P. Fierlinger, D. Fudenberg, R. Gornea, K. Graham, G. Gratta, C. Hall, E. V. Hansen, J. Hoessl, P. Hufschmidt, M. Hughes, A. Jamil, M. J. Jewell, A. Johnson, S. Johnston, A. Karelin, L. J. Kaufman, T. Koffas, S. Kravitz, R. Krücken, A. Kuchenkov, K. S. Kumar, Y. Lan, D. S. Leonard, G. S. Li, S. Li, C. Licciardi, Y. H. Lin, R. MacLellan, T. Michel, B. Mong, D. Moore, K. Murray, R. Nelson, O. Njoya, A. Odian, I. Ostrovskiy, A. Piepke, A. Pocar, F. Retière, P. C. Rowson, J. J. Russell, S. Schmidt, A. Schubert, D. Sinclair, V. Stekhanov, M. Tarka, T. Tolba, R. Tsang, P. Vogel, J.-L. Vuilleumier, M. Wagenpfeil, A. Waite, T. Walton, M. Weber, L. J. Wen, U. Wichoski, G. Wrede, L. Yang, Y.-R. Yen, O. Y. Zeldovich, J. Zettlemoyer, T. Ziegler, Search for Neutrinoless Double- β Decay with the Upgraded EXO-200 Detector. *Phys. Rev. Lett.* **120**, 072701 (2018). [doi:10.1103/PhysRevLett.120.072701](https://doi.org/10.1103/PhysRevLett.120.072701) [Medline](https://arxiv.org/abs/1805.02550)
31. K. H. Ackermann, M. Agostini, M. Allardt, M. Altmann, E. Andreotti, A. M. Bakalyarov, M. Balata, I. Barabanov, M. Barnabé Heider, N. Barros, L. Baudis, C. Bauer, N. Becerici-Schmidt, E. Bellotti, S. Belogurov, S. T. Belyaev, G. Benato, A. Bettini, L. Bezrukov, T. Bode, V. Brudanin, R. Brugnera, D. Budjáš, A. Caldwell, C. Cattadori, A. Chernogorov, O. Chkvetse, F. Cossavella, A. D'Andragora, E. V. Demidova, A. Denisov, A. di Vacri, A. Domula, V. Egorov, R. Falkenstein, A. Ferella, K. Freund, F. Froberg, N. Frodyma, A. Gangapshev, A. Garfagnini, J. Gasparro, S. Gazzana, R. Gonzalez de Orduna, P. Grabmayr, V. Gurentsov, K. Gusev, K. K. Guthikonda, W. Hampel, A. Hegai, M. Heisel, S. Hemmer, G. Heusser, W. Hofmann, M. Hult, L. V. Inzhchik, L. Ioannucci, J. Janicskó Csáthy, J. Jochum, M. Junker, R. Kankanyan, S. Kianovsky, T. Kihm, J. Kiko, I. V. Kirpichnikov, A. Kirsch, A. Klimenko, M. Knapp, K. T. Knöpfle, O. Kochetov, V. N. Kornoukhov, K. Kröninger, V. Kusminov, M. Laubenstein, A. Lazzaro, V. I. Lebedev, B. Lehnert, D. Lenz, H. Liao, M. Lindner, I. Lippi, J. Liu, X. Liu, A. Lubashevskiy, B. Lubsandorzhiev, A. A. Machado, B. Majorovits, W. Maneschg, G. Marissens, S. Mayer, G. Meierhofer, I. Nemchenok, L. Niedermeier, S. Nisi, J. Oehm, C. O'Shaughnessy, L. Pandola, P. Peiffer, K. Pelczar, A. Pullia, S. Riboldi, F. Ritter, C. Rossi Alvarez, C. Sada, M. Salathe, C. Schmitt, S. Schönert, J. Schreiner, J. Schubert, O. Schulz, U. Schwan, B. Schwingenheuer, H. Seitz, E. Shevchik, M. Shirchenko, H. Simgen, A. Smolnikov, L. Stanco, F. Stelzer, H. Strecker, M. Tarka, U. Trunk, C. A. Ur, A. A. Vasenko, S. Vogt, O. Volynets, K. von Sturm, V. Wagner, M. Walter, A. Wegmann, M. Wojcik, E. Yanovich, P. Zavarise, I. Zhitnikov, S. V. Zhukov, D. Zinatulina, K. Zuber, G. Zuzel, The Gerda experiment for the search of $0\nu\beta\beta$ decay in ^{76}Ge . *Eur. Phys. J. C* **73**, 2330 (2013). [doi:10.1140/epjc/s10052-013-2330-0](https://doi.org/10.1140/epjc/s10052-013-2330-0)
32. W. Maneschg, M. Laubenstein, D. Budjáš, W. Hampel, G. Heusser, K. T. Knöpfle, B. Schwingenheuer, H. Simgen, Measurements of extremely low radioactivity levels in stainless steel for GERDA. *Nucl. Instrum. Methods A* **593**, 448–453 (2008). [doi:10.1016/j.nima.2008.05.036](https://doi.org/10.1016/j.nima.2008.05.036)

33. I. Barabanov, L. Bezrukov, E. Demidova, V. Gurentsov, S. Kianovsky, K. T. Knöpfle, V. Kornoukhov, B. Schwingenheuer, A. Vasenko, Shielding of the GERDA experiment against external gamma background. *Nucl. Instrum. Methods A* **606**, 790–794 (2009). [doi:10.1016/j.nima.2009.04.006](https://doi.org/10.1016/j.nima.2009.04.006)
34. K. Freund, R. Falkenstein, P. Grabmayr, A. Hegai, J. Jochum, M. Knapp, B. Lubsandorzhiev, F. Ritter, C. Schmitt, A.-K. Schütz, I. Jitnikov, E. Shevchik, M. Shirchenko, D. Zinatulina, The performance of the Muon Veto of the Gerda experiment. *Eur. Phys. J. C* **76**, 298 (2016). [doi:10.1140/epjc/s10052-016-4140-7](https://doi.org/10.1140/epjc/s10052-016-4140-7)
35. M. Günther, J. Hellmig, G. Heusser, M. Hirsch, H. V. Klapdor-Kleingrothaus, B. Maier, H. Päs, F. Petry, Y. Ramachers, H. Strecker, M. Völlinger, A. Balysh, S. T. Belyaev, A. Demehin, A. Gurov, I. Kondratenko, D. Kotelnikov, V. I. Lebedev, A. Müller, Heidelberg-Moscow $\beta\beta$ experiment with ^{76}Ge : Full setup with five detectors. *Phys. Rev. D* **55**, 54–67 (1997). [doi:10.1103/PhysRevD.55.54](https://doi.org/10.1103/PhysRevD.55.54)
36. C. E. Aalseth, F. T. Avignone, R. L. Brodzinski, S. Cebrian, E. Garcia, D. Gonzalez, W. K. Hensley, I. G. Irastorza, I. V. Kirpichnikov, A. A. Klimenko, H. S. Miley, A. Morales, J. Morales, A. O. de Solorzano, S. B. Osetrov, V. S. Pogosov, J. Puimiedon, J. H. Reeves, M. L. Sarsa, A. A. Smolnikov, A. S. Starostin, A. G. Tamanyan, A. A. Vasenko, S. I. Vasiliev, J. A. Villar, IGEX ^{76}Ge neutrinoless double-beta decay experiment: Prospects for next generation experiments. *Phys. Rev. D* **65**, 092007 (2002). [doi:10.1103/PhysRevD.65.092007](https://doi.org/10.1103/PhysRevD.65.092007)
37. M. Agostini, M. Allardt, E. Andreotti, A. M. Bakalyarov, M. Balata, I. Barabanov, M. Barnabé-Heider, N. Barros, L. Baudis, C. Bauer, N. Becerici-Schmidt, E. Bellotti, S. Belogurov, S. T. Belyaev, G. Benato, A. Bettini, L. Bezrukov, T. Bode, V. Brudanin, R. Brugnera, D. Budjáš, A. Caldwell, C. Cattadori, A. Chernogorov, F. Cossavella, E. V. Demidova, A. Domula, V. Egorov, R. Falkenstein, A. Ferella, K. Freund, N. Frodyma, A. Gangapshev, A. Garfagnini, C. Gotti, P. Grabmayr, V. Gurentsov, K. Gusev, K. K. Guthikonda, W. Hampel, A. Hegai, M. Heisel, S. Hemmer, G. Heusser, W. Hofmann, M. Hult, L. V. Inzhechik, L. Ioannucci, J. Janicskó Csáthy, J. Jochum, M. Junker, T. Kihm, I. V. Kirpichnikov, A. Kirsch, A. Klimenko, K. T. Knöpfle, O. Kochetov, V. N. Kornoukhov, V. V. Kuzminov, M. Laubenstein, A. Lazzaro, V. I. Lebedev, B. Lehnert, H. Y. Liao, M. Lindner, I. Lippi, X. Liu, A. Lubashevskiy, B. Lubsandorzhiev, G. Lutter, C. Macolino, A. A. Machado, B. Majorovits, W. Maneschg, M. Misiaszek, I. Nemchenok, S. Nisi, C. O'Shaughnessy, L. Pandola, K. Pelczar, G. Pessina, A. Pullia, S. Riboldi, N. Romyantseva, C. Sada, M. Salathe, C. Schmitt, J. Schreiner, O. Schulz, B. Schwingenheuer, S. Schönert, E. Shevchik, M. Shirchenko, H. Simgen, A. Smolnikov, L. Stanco, H. Strecker, M. Tarka, C. A. Ur, A. A. Vasenko, O. Volynets, K. von Sturm, V. Wagner, M. Walter, A. Wegmann, T. Wester, M. Wojcik, E. Yanovich, P. Zavarise, I. Zhitnikov, S. V. Zhukov, D. Zinatulina, K. Zuber, G. Zuzel, Pulse shape discrimination for Gerda Phase I data. *Eur. Phys. J. C* **73**, 2583 (2013). [doi:10.1140/epjc/s10052-013-2583-7](https://doi.org/10.1140/epjc/s10052-013-2583-7)
38. M. Agostini, M. Allardt, E. Andreotti, A. M. Bakalyarov, M. Balata, I. Barabanov, N. Barros, L. Baudis, C. Bauer, N. Becerici-Schmidt, E. Bellotti, S. Belogurov, S. T. Belyaev, G. Benato, A. Bettini, L. Bezrukov, T. Bode, D. Borowicz, V. Brudanin, R. Brugnera, D. Budjáš, A. Caldwell, C. Cattadori, A. Chernogorov, V. D'Andrea, E. V. Demidova, A. Domula, V. Egorov, R. Falkenstein, K. Freund, N. Frodyma, A. Gangapshev, A. Garfagnini, C. Gotti, P. Grabmayr, V. Gurentsov, K. Gusev, A. Hegai, M. Heisel, S. Hemmer, G. Heusser, W. Hofmann, M. Hult, L. V. Inzhechik, L. Ioannucci, J. Janicskó Csáthy, J. Jochum, M. Junker, V. Kazalov, T. Kihm, I. V. Kirpichnikov, A. Kirsch, A. Klimenko, K. T. Knöpfle, O. Kochetov, V. N. Kornoukhov, V. V. Kuzminov, M. Laubenstein, A. Lazzaro, V. I. Lebedev, B. Lehnert, H. Y. Liao, M. Lindner, I. Lippi, A. Lubashevskiy, B. Lubsandorzhiev, G. Lutter, C. Macolino, B. Majorovits, W. Maneschg, M. Misiaszek, I. Nemchenok, S. Nisi, C. O'Shaughnessy, D. Palioselitis, L. Pandola, K. Pelczar, G. Pessina, A. Pullia, S. Riboldi, N. Romyantseva, C. Sada, M. Salathe, C. Schmitt, J. Schreiner, O. Schulz, A.-K. Schütz, B. Schwingenheuer, S. Schönert, E. Shevchik, M. Shirchenko, H. Simgen, A. Smolnikov, L. Stanco, H. Strecker, C. A. Ur, L. Vanhoefer, A. A. Vasenko, K. von Sturm, V. Wagner, M. Walter, A. Wegmann, T. Wester, H. Wilsenach, M. Wojcik, E. Yanovich, P. Zavarise, I. Zhitnikov, S. V. Zhukov, D. Zinatulina, K. Zuber, G. Zuzel, Production, characterization and operation of ^{76}Ge enriched BEGe detectors in GERDA. *Eur. Phys. J. C* **75**, 39 (2015). [doi:10.1140/epjc/s10052-014-3253-0](https://doi.org/10.1140/epjc/s10052-014-3253-0)
39. S. Riboldi et al., in *Proceedings, 4th International Conference on Advancements in Nuclear Instrumentation Measurement Methods and their Applications (ANIMMA 2015)*, 7465549 (2015). [doi:10.1109/ANIMMA.2015.7465549](https://doi.org/10.1109/ANIMMA.2015.7465549)
40. M. Agostini, M. Barnabé-Heider, D. Budjáš, C. Cattadori, A. Gangapshev, K. Gusev, M. Heisel, M. Junker, A. Klimenko, A. Lubashevskiy, K. Pelczar, S. Schönert, A. Smolnikov, G. Zuzel, LArGe: Active background suppression using argon scintillation for the Gerda $0\nu\beta\beta$ experiment. *Eur. Phys. J. C* **75**, 506 (2015). [doi:10.1140/epjc/s10052-015-3681-5](https://doi.org/10.1140/epjc/s10052-015-3681-5)
41. J. Janicskó Csáthy, H. Aghaei Khozani, A. Caldwell, X. Liu, B. Majorovits, Development of an anti-Compton veto for HPGe detectors operated in liquid argon using silicon photo-multipliers. *Nucl. Instrum. Methods A* **654**, 225–232 (2011). [doi:10.1016/j.nima.2011.05.070](https://doi.org/10.1016/j.nima.2011.05.070)
42. A. Lubashevskiy, M. Agostini, D. Budjáš, A. Gangapshev, K. Gusev, M. Heisel, A. Klimenko, A. Lazzaro, B. Lehnert, K. Pelczar, S. Schönert, A. Smolnikov, M. Walter, G. Zuzel, Mitigation of $^{42}\text{Ar}/^{42}\text{K}$ background for the GERDA Phase II experiment. *Eur. Phys. J. C* **78**, 15 (2018). [doi:10.1140/epjc/s10052-017-5499-9](https://doi.org/10.1140/epjc/s10052-017-5499-9)
43. M. Agostini, J. A. Detwiler, P. Finnerty, K. Kröninger, D. Lenz, J. Liu, M. G. Marino, R. Martin, K. D. Nguyen, L. Pandola, A. G. Schubert, O. Volynets, P. Zavarise, The MGDO software library for data analysis in Ge neutrinoless double-beta decay experiments. *J. Phys. Conf. Ser.* **375**, 042027 (2012). [doi:10.1088/1742-6596/375/1/042027](https://doi.org/10.1088/1742-6596/375/1/042027)
44. M. Agostini et al., GELATIO: A general framework for modular digital analysis of high-purity Ge detector signals. *J. Instrum.* **6**, P08013 (2011). [doi:10.1088/1748-0221/6/08/P08013](https://doi.org/10.1088/1748-0221/6/08/P08013)
45. M. Agostini, L. Pandola, P. Zavarise, Off-line data processing and analysis for the GERDA experiment. *J. Phys. Conf. Ser.* **368**, 012047 (2012). [doi:10.1088/1742-6596/368/1/012047](https://doi.org/10.1088/1742-6596/368/1/012047)
46. M. Agostini, M. Allardt, A. M. Bakalyarov, M. Balata, I. Barabanov, N. Barros, L. Baudis, C. Bauer, N. Becerici-Schmidt, E. Bellotti, S. Belogurov, S. T. Belyaev, G. Benato, A. Bettini, L. Bezrukov, T. Bode, D. Borowicz, V. Brudanin, R. Brugnera, D. Budjáš, A. Caldwell, C. Cattadori, A. Chernogorov, V. D'Andrea, E. V. Demidova, A. Vacri, A. Domula, E. Doroshkevich, V. Egorov, R. Falkenstein, O. Fedorova, K. Freund, N. Frodyma, A. Gangapshev, A. Garfagnini, P. Grabmayr, V. Gurentsov, K. Gusev, A. Hegai, M. Heisel, S. Hemmer, G. Heusser, W. Hofmann, M. Hult, L. V. Inzhechik, J. Janicskó Csáthy, J. Jochum, M. Junker, V. Kazalov, T. Kihm, I. V. Kirpichnikov, A. Kirsch, A. Klimenko, K. T. Knöpfle, O. Kochetov, V. N. Kornoukhov, V. V. Kuzminov, M. Laubenstein, A. Lazzaro, V. I. Lebedev, B. Lehnert, H. Y. Liao, M. Lindner, I. Lippi, A. Lubashevskiy, B. Lubsandorzhiev, G. Lutter, C. Macolino, B. Majorovits, W. Maneschg, E. Medinaceli, M. Misiaszek, P. Moseev, I. Nemchenok, D. Palioselitis, K. Panas, L. Pandola, K. Pelczar, A. Pullia, S. Riboldi, N. Romyantseva, C. Sada, M. Salathe, C. Schmitt, B. Schneider, S. Schönert, J. Schreiner, A.-K. Schütz, O. Schulz, B. Schwingenheuer, O. Selivanenko, M. Shirchenko, H. Simgen, A. Smolnikov, L. Stanco, M. Stepaniuk, C. A. Ur, L. Vanhoefer, A. A. Vasenko, A. Veresnikova, K. von Sturm, V. Wagner, M. Walter, A. Wegmann, T. Wester, H. Wilsenach, M. Wojcik, E. Yanovich, P. Zavarise, I. Zhitnikov, S. V. Zhukov, D. Zinatulina, K. Zuber, G. Zuzel, Improvement of the energy resolution via an optimized digital signal processing in GERDA Phase I. *Eur. Phys. J. C* **75**, 255 (2015). [doi:10.1140/epjc/s10052-015-3409-6](https://doi.org/10.1140/epjc/s10052-015-3409-6)
47. L. Baudis, A. D. Ferella, F. Froborg, M. Tarka, Monte Carlo studies and optimization for the calibration system of the Gerda experiment. *Nucl. Instrum. Methods A* **729**, 557–564 (2013). [doi:10.1016/j.nima.2013.08.003](https://doi.org/10.1016/j.nima.2013.08.003)
48. L. Baudis et al., Production and characterization of ^{228}Th calibration sources with low neutron emission for GERDA. *J. Instrum.* **10**, P12005 (2015). [doi:10.1088/1748-0221/10/12/P12005](https://doi.org/10.1088/1748-0221/10/12/P12005)
49. M. Agostini, M. Allardt, A. M. Bakalyarov, M. Balata, I. Barabanov, N. Barros, L. Baudis, C. Bauer, N. Becerici-Schmidt, E. Bellotti, S. Belogurov, S. T. Belyaev, G. Benato, A. Bettini, L. Bezrukov, T. Bode, D. Borowicz, V. Brudanin, R. Brugnera, D. Budjáš, A. Caldwell, C. Cattadori, A. Chernogorov, V. D'Andrea, E. V. Demidova, A. di Vacri, A. Domula, E. Doroshkevich, V. Egorov, R. Falkenstein, O. Fedorova, K. Freund, N. Frodyma, A. Gangapshev, A. Garfagnini, P. Grabmayr, V. Gurentsov, K. Gusev, A. Hegai, M. Heisel, S. Hemmer, G. Heusser, W. Hofmann,

- M. Hult, L. V. Inzhechik, J. J. Csáthy, J. Jochum, M. Junker, V. Kazalov, T. Kihm, I. V. Kirpichnikov, A. Kirsch, A. Klimenko, K. T. Knöpfle, O. Kochetov, V. N. Kornoukhov, V. V. Kuzminov, M. Laubenstein, A. Lazzaro, V. I. Lebedev, B. Lehnert, H. Y. Liao, M. Lindner, I. Lippi, A. Lubashevskiy, B. Lubsandorzhiev, G. Lutter, C. Macolino, B. Majorovits, W. Maneschg, E. Medinaceli, M. Misiaszek, P. Moseev, I. Nemchenok, D. Palioselitis, K. Panas, L. Pandola, K. Pelczar, A. Pullia, S. Riboldi, N. Rumyantseva, C. Sada, M. Salathe, C. Schmitt, B. Schneider, S. Schöner, J. Schreiner, A.-K. Schütz, O. Schulz, B. Schwingenheuer, O. Selivanenko, M. Shirchenko, H. Simgen, A. Smolnikov, L. Stanco, M. Stepaniuk, C. A. Ur, L. Vanhoefer, A. A. Vasenko, A. Veresnikova, K. von Sturm, V. Wagner, M. Walter, A. Wegmann, T. Wester, H. Wilsenach, M. Wojcik, E. Yanovich, P. Zavarise, I. Zhitnikov, S. V. Zhukov, D. Zinatulina, K. Zuber, G. Zuzel, Results on $\beta\beta$ decay with emission of two neutrinos or Majorons in ^{76}Ge from GERDA Phase I. *Eur. Phys. J. C* **75**, 416 (2015). doi:10.1140/epjc/s10052-015-3627-y
50. J. Kotila, F. Iachello, Phase-space factors for double- β decay. *Phys. Rev. C* **85**, 034316 (2012). doi:10.1103/PhysRevC.85.034316
51. D. Štefánik, R. Dvornický, F. Šimkovic, P. Vogel, Reexamining the light neutrino exchange mechanism of the $0\nu\beta\beta$ decay with left- and right-handed leptonic and hadronic currents. *Phys. Rev. C* **92**, 055502 (2015). doi:10.1103/PhysRevC.92.055502
52. J. Menéndez, A. Poves, E. Caurier, F. Nowacki, Disassembling the nuclear matrix elements of the neutrinoless $\beta\beta$ decay. *Nucl. Phys. A* **818**, 139–151 (2009). doi:10.1016/j.nuclphysa.2008.12.005
53. M. Horoi, A. Neacsu, Shell model predictions for ^{124}Sn double- β decay. *Phys. Rev. C* **93**, 024308 (2016). doi:10.1103/PhysRevC.93.024308
54. J. M. Yao, L. S. Song, K. Hagino, P. Ring, J. Meng, Systematic study of nuclear matrix elements in neutrinoless double- β decay with a beyond-mean-field covariant density functional theory. *Phys. Rev. C* **91**, 024316 (2015). doi:10.1103/PhysRevC.91.024316
55. N. López Vaquero, T. R. Rodríguez, J. L. Egido, Shape and pairing fluctuation effects on neutrinoless double β decay nuclear matrix elements. *Phys. Rev. Lett.* **111**, 142501 (2013). doi:10.1103/PhysRevLett.111.142501 Medline
56. J. Barea, J. Kotila, F. Iachello, $0\nu\beta\beta$ and $2\nu\beta\beta$ nuclear matrix elements in the interacting boson model with isospin restoration. *Phys. Rev. C* **91**, 034304 (2015). doi:10.1103/PhysRevC.91.034304
57. J. Hyvärinen, J. Suhonen, Nuclear matrix elements for $0\nu\beta\beta$ decays with light or heavy Majorana-neutrino exchange. *Phys. Rev. C* **91**, 024613 (2015). doi:10.1103/PhysRevC.91.024613
58. M. T. Mustonen, J. Engel, Large-scale calculations of the double- β decay of ^{76}Ge , ^{130}Te , ^{136}Xe , and ^{150}Nd in the deformed self-consistent Skyrme quasiparticle random-phase approximation. *Phys. Rev. C* **87**, 064302 (2013). doi:10.1103/PhysRevC.87.064302
59. F. Šimkovic, V. Rodin, A. Faessler, P. Vogel, $0\nu\beta\beta$ and $2\nu\beta\beta$ nuclear matrix elements, quasiparticle random-phase approximation, and isospin symmetry restoration. *Phys. Rev. C* **87**, 045501 (2013). doi:10.1103/PhysRevC.87.045501

ACKNOWLEDGMENTS

The GERDA collaboration thanks the directors and the staff of the LNGS for their continuous strong support of the GERDA experiment. **Funding:** The GERDA experiment is supported financially by the German Federal Ministry for Education and Research (BMBF), the German Research Foundation (DFG) via the Excellence Cluster Universe and the SFB1258, the Italian Istituto Nazionale di Fisica Nucleare (INFN), the Max Planck Society (MPG), the Polish National Science Center (NCN) under grant UMO-2016/21/B/ST2/01094, the Foundation for Polish Science (TEAM/2016-2/2017), the Russian Foundation for Basic Research (RFBR) and the Swiss National Science Foundation (SNF). The institutions acknowledge also internal financial support. This project has received funding or support from the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie Grant Agreements No. 690575 and No. 674896, respectively. **Author contributions:** All authors contributed to the publication, being differently involved in the design and construction of the detector system, in its operation, and

in the acquisition and analysis of data. All authors approved the final version of the manuscript. In line with collaboration policy, the authors are listed alphabetically. **Competing interests:** The authors declare no competing financial interests. **Data and materials availability:** All data generated during this analysis and shown in Fig. 2, Fig. 3, and figs. S3 to S12 are available as png, pdf and root/txt files from the GERDA repository (26) at Zenodo. For further information contact the GERDA Collaboration (gerda-eb@mpi-hd.mpg.de).

SUPPLEMENTARY MATERIALS

science.sciencemag.org/cgi/content/full/science.aav8613/DC1

Materials and Methods

Supplementary Text

Figs. S1 to S12

Tables S1 to S7

References (31–59)

27 November 2018; accepted 20 August 2019

Published online 5 September 2019

10.1126/science.aav8613

Table 1. Comparison of present and prior experiments. Lower half-life limits $L(T_{1/2})$ and sensitivities $S(T_{1/2})$, both at 90% C.L., reported by recent $0\nu\beta\beta$ decay searches with indicated deployed isotope masses M_i and FWHM energy resolutions. Sensitivities $S(T_{1/2})$ have been converted into upper limits of effective Majorana masses $m_{\beta\beta}$ using the nuclear matrix elements quoted in (20).

Experiment	Isotope	M_i (kmol)	FWHM (keV)	$L(T_{1/2})$ (10^{25} yr)	$S(T_{1/2})$ (10^{25} yr)	$m_{\beta\beta}$ (meV)
GERDA (this work)	^{76}Ge	0.41	3.3	9	11	104-228
MAJORANA (27)	^{76}Ge	0.34	2.5	2.7	4.8	157-346
CUPID-0 (28)	^{82}Se	0.063	23	0.24	0.23	394-810
CUORE (29)	^{130}Te	1.59	7.4	1.5	0.7	162-757
EXO-200 (30)	^{136}Xe	1.04	71	1.8	3.7	93-287
KamLAND-Zen (21)	^{136}Xe	2.52	270	10.7	5.6	76-234
Combined						66-155

Fig. 1. The concept of active background suppression. Gerda searches for the $0\nu\beta\beta$ decay $^{76}\text{Ge} \rightarrow ^{76}\text{Se} + 2e^{-}$, $Q_{\beta\beta} = 2039$ keV, with high-purity Ge detectors enriched in ^{76}Ge that are operated in liquid argon (LAr). (i), Artist's view of the $0\nu\beta\beta$ decay of a nucleus by an emission of two electrons (β particles); below, 3 BEGe detectors, (ii) to (iv), out of the 40 Ge detectors of the Gerda detector array (table S1 and fig. S2) immersed in LAr (bluish cylinder). Events from $0\nu\beta\beta$ decays would deposit energy $Q_{\beta\beta}$ within a few cubic millimeters in a single detector (ii). Events with coincident LAr scintillation light or with multiple interactions in the Ge detector, e.g., from Compton scattering (iii), are classified as background events. The special detector design with a small readout electrode (fig. S1) enhances drift time differences between different trajectories (black dashed lines) of the charges (holes) generated by the energy depositions. The color code (see fig. S1 for color bar) indicates the electrical signal strength at the respective location. Hence single- and multi-site events can be identified efficiently by the time profile of their electronic signal. Similarly, α decays at the readout electrode show unique signal characteristics (iv).

Fig. 2. Gerda Phase II energy spectra (53.9 kg · yr). Enriched coaxial and BEGe data are displayed in a combined spectrum after indicated cuts. Main contributions to the spectra are labeled. The insets display the analysis window for both coaxial and BEGe detectors separately including the background rates (solid blue lines). No event reconstructs within $Q_{\beta\beta} \pm 2\sigma$. The dashed blue curves depict the 90% C.L. limit for a $0\nu\beta\beta$ signal of $T_{1/2}^{0\nu} = 0.9 \cdot 10^{26}$ yr derived from the likelihood analysis of all Gerda data sets.

Fig. 3. Constraints of the parameter space for $m_{\beta\beta}$ in the scenario of 3 light Majorana neutrinos. Constraints are shown, left to right, as function of the lightest neutrino mass m_{light} , the sum of neutrino masses Σ , and the effective neutrino mass m_{β} . Contours follow from a scan of the Majorana phases with the central oscillation parameters from NuFIT 4.0 (22). The blue horizontal band shows the upper limits on $m_{\beta\beta}$ obtained by Gerda, the grey band those from combining sensitivities of all leading experiments in the field (see Table 1). Vertical lines denote $\Sigma = 0.12$ eV and $\Sigma = 0.66$ eV, a stringent limit from cosmology (24) and an extended model bound (7), as well as $m_{\beta} = 0.23$ eV, the 5 yr sensitivity of the KATRIN experiment (23). Hatchures tag the excluded parameter space.