Cite as: D. Kong *et al.*, *Science*
10.1126/science.abb4129 (2020).

Direct reversible decarboxylation from stable organic acids in dimethylformamide solution

Duanyang Kong*, Patrick J. Moon*, Erica K. J. Lui, Odey Bsharat, Rylan J. Lundgren†

Department of Chemistry, University of Alberta, Edmonton, Alberta T6G 2G2, Canada.

*These authors contributed equally to this work. †Corresponding author. Email: rylan.lundgren@ualberta.ca

Many classical and emerging methodologies in organic chemistry rely on CO₂ extrusion to generate reactive intermediates for bond-forming events. Synthetic reactions that involve the microscopic reverse, the carboxylation of reactive intermediates, have conventionally been undertaken using very different conditions. We report that chemically stable C(sp³) carboxylates, such as arylacetic acids and malonate half-esters, undergo uncatalyzed reversible decarboxylation in dimethylformamide solution.

Decarboxylation/carboxylation occurs with substrates resistant to protodecarboxylation by Brønsted acids under otherwise identical conditions. Isotopically labeled carboxylic acids can be prepared in high chemical and isotopic yield by simply supplying an atmosphere of ¹³CO₂ to carboxylate salts in polar aprotic solvents. An understanding of carboxylate reactivity in solution enables conditions for the trapping of aldehydes, ketones, and α,β-unsaturated esters.

Decarboxylation is a fundamental step in biochemical processes and synthetic organic chemistry. Fermentation, respiration, and the biosynthesis of many secondary metabolites involve the loss of CO₂ from organic acids (1, 2). Decarboxylase enzymes accelerate these reactions by stabilizing developing intermediates (typically carbanions) and promoting CO₂ diffusion from the active site, thereby enabling otherwise unfeasible decarboxylations to occur under physiological conditions (Fig. 1A) (3, 4). Acid substrates lacking strong anion-stabilizing groups adjacent to the reactive carbon center have been construed to be inert toward spontaneous decarboxylation without resorting to thermolysis conditions (Fig. 1B) (5). As a result, synthetic reactions driven by decarboxylation are often carried out using high reaction temperatures (6), added oxidizing agents (7, 8), or prior stoichiometric chemical modification of the carboxylate unit (9–12).

Carboxylation reactions, the microscopic reverse of decarboxylations, are equally valuable processes in biology and synthetic chemistry. Despite the possibility of a shared reaction pathway, the biochemical machinery that promotes carboxylation in CO₂ fixation pathways operates with a distinct set of substrates and enzymes from those that promote decarboxylation in all but a few cases (13–17). Similarly, synthetic techniques that generate carboxylic acid derivatives from CO₂ have tended to apply strongly nucleophilic organometallics and/or in-situ stoichiometric (electro)chemical substrate reduction (18, 19).

The potential for the reversibility of decarboxylation/carboxylation mechanisms is largely ignored in reports of synthetic methodologies that rely on these elementary steps. Reports of direct non-enzymatic reversible CO₂-exchange of

carboxylic acids are restricted to specialized substrate/mediator pairs (20, 21). Exchange of carboxylate groups in simple aliphatic acids with CO₂ has been documented, but requires heating of neat substrates at 280–400°C (22, 23). Nonetheless, in the course of our studies on catalytic decarboxylative cross-coupling reactions (24, 25), we questioned whether the apparent stability of organic carboxylates could arise from reversible decarboxylation/carboxylation events in solution. Supporting this hypothesis, we observed that certain simple organic acids that are stable toward protodecarboxylation in solution undergo spontaneous incorporation of ¹³CO₂ when this heavier isotope is supplied at atmospheric pressure (Fig. 1C).

The potassium salt of arylacetic acid **1** exemplifies the reversible decarboxylation/carboxylation behavior of otherwise chemically stable carboxylic acids. A 0.1 M solution of **1** in dimethylformamide (DMF) at 20°C underwent CO₂ exchange when placed under an atmosphere of ¹³CO₂. In a reaction where approximately six equivalents of ¹³CO₂ were supplied (13 mL of CO₂ at ~1 atm, dissolved [CO₂] = 0.20 M), equilibrium between ¹²C and ¹³C was achieved in 15 hours (Fig. 1C, red trace). Quantitative recovery of carboxylate **1** with 83% ¹³C-enrichment was possible by acid/base extractive workup. Under similar conditions at 20°C with five equivalents of a weak Brønsted acid (MeOH) no protodecarboxylation of **1** was observed (Fig. 1C, black trace). These results demonstrate that capture of the putative nucleophilic intermediate generated from **1** with dissolved CO₂ is significantly more favorable than protonation. The process tolerates up to 0.01 M H₂O and does not require rigorous exclusion of air (see fig. S2). During the review of this work, isotopic exchange of carboxylate

groups in cesium arylacetic acid salts with ^{14}C -, ^{13}C -, and ^{11}C -labeled CO_2 in DMSO at 80–190°C was reported (26).

The counter-cation of the carboxylate salt impacts carboxylate exchange reactivity (Fig. 1D). The carboxylic acid of **1** manifested no evident $^{13}\text{CO}_2$ exchange or protodecarboxylation in DMF at 70°C. Li^+ and Na^+ salts of **1** reacted more slowly, while the Cs^+ salt reacted more quickly. Divalent metal salts of **1** (Zn^{2+} or Cu^{2+}) were inert and the addition of M^{2+}Cl_2 salts completely inhibited the reaction (MgCl_2 , CaCl_2 , MnCl_2 , see fig. S3). The use of polar aprotic solvents (DMF, DMA, DMSO; dielectric constant $\epsilon > 30$) is essential for the transformation: reactions conducted in THF, DCE, or water resulted in recovery of unlabeled **1** at 20°C. The addition of 18-crown-6 (18-C-6) led to an approximate two-fold rate enhancement of carboxylate exchange (see fig. S4) (27). The free acid underwent carboxylate exchange when 1.5 equivalents of K_2CO_3 and 18-C-6 were added (>90% yield and ^{13}C incorporation in 19 hours). Collectively, these observations suggest that the generation of a solvent-separated ion pair leads to enhanced decarboxylative reactivity.

Reversible decarboxylation occurred for an array of carboxylate containing molecules that contain adjacent aryl, carbonyl, cyano, or sulfonyl groups, including valuable synthetic precursors, drug molecules, and amino acid derivatives (Fig. 2). The incorporation of $^{13}\text{CO}_2$ and product recovery remained high (>80%) across several substrate classes. The degree of incorporation is largely a function of the amount of $^{13}\text{CO}_2$ supplied: >95% enrichment can be obtained when ~50 equivalents is provided (see fig. S5). For successful cases, this carboxylate exchange process compares favorably in terms of operational simplicity to current state-of-the-art methods to prepare $\text{C}(\text{sp}^3)\text{-}^{13/14}\text{CO}_2$ labeled carboxylic acid-derivatives sought after in (pre)clinical absorption, distribution, metabolism, and excretion (ADME) studies (28). Reported approaches require either chemical activation-decarboxylation-metalation-carboxylation sequences mediated by transition metals (29–31), indirect nucleophilic substitution reactions with labeled cyanide followed by hydrolysis (32), or introduction of labeled carbon monoxide in place of CO_2 (33, 34). The direct exchange of $\text{C}(\text{sp}^2)$ -carboxylate groups catalyzed by transition metals has been demonstrated; however, reactivity is restricted to nitro- or sulfonyl-containing arenes or 2-heteroatom substituted electron-rich heterocycles at high temperature ($\geq 150^\circ\text{C}$) (35).

(Hetero)arylacetic acid salts with anion-stabilizing groups underwent exchange at moderate temperatures (Fig. 2, **1–4**, **9**, **10**, **11–15** at 20 to 80°C), whereas arylacetates with strongly electron-donating OMe or NMe₂ groups required higher temperatures (**17–20** at 100 to 130°C) and benefitted from the addition of 18-C-6. The simplicity of the process enabled broad functional group compatibility, including tolerance to boronic esters (**6**), aryl halides (I, Br, Cl, F; **4**, **7**, **8**,

10), ketones (**11**), aldehydes (**12**), esters (**14**), amides (**13**), sulfonyls (**15**), and potentially reactive heterocycles (chromenone **25**, NH-indole **26**, pyridines **27**, **29**, pyrimidine **28**, isoxazole **30**, thiophene **31**). Alkyl and aryl substitution adjacent to the carboxylate was tolerated, including examples of trisubstituted, non-enolizable arylacetic acid salts (**23**, **24**). Other classes of potassium carboxylates that underwent reversible decarboxylation include malonate half-esters (**32–35**), keto acids (**36**), β -carboxysulfonyls (**37**, **38**), cyanoacetates (**39**), and carboxylactams (**40**). Alkene and terminal alkyne functional groups did not interfere with the process (**34**, **35**). Potassium malonates underwent CO_2 exchange at higher temperature (135°C) to give a mixture of mono- and doubly-labeled product along with ^{13}C -enriched monoacid (**41**, **42**).

Carboxylate exchange could also be used to directly prepare isotopically labeled drug molecules, including arylacetic acid salts and propionate NSAIDs of varying complexity (**43–52**, Fig. 2). Pharmaceuticals featuring amide or ester groups were obtained via derivatization of the acid group (Zolpidem **53**, Aprofene **55**) or could be prepared according to established literature protocols (Propiverine **54**, Netupitant **56**, Repaglinide **57**). Consistent with the generation of a carbanion, racemization of enantiopure Naproxen (**46**) was observed (fig. S6). Reversible decarboxylation may explain reports of aryl propionate racemization required for kinetic resolution manufacturing processes (36, 37). Simple alkyl carboxylates did not undergo CO_2 exchange; however isotopically labeled products of this class can be readily obtained by carboxylate exchange/desulfonylation reactions of β -sulfonyl acids or exchange/decarboxylation sequences of malonic acids in three steps (**58–60**). The facile generation of ^{13}C -diphenylmethylidene glycine at room temperature (**61** 93% incorporation, 76% yield) serves as a starting point for the synthesis of other labeled amino acids (38).

The reversible CO_2 exchange process likely involves the formation of a carbon nucleophile either from direct decarboxylation, or potentially in the case of enolizable substrates, through an enolate intermediate. The reaction rates and required temperatures for $^{12}\text{CO}_2$ / $^{13}\text{CO}_2$ interconversion correlate with the substrate's capacity to stabilize negative charge and not with oxidation potential (compare **1**, **14**, **16**, and **17**). The addition of radical inhibitors (TEMPO, BHT) had no impact on the decarboxylative reactivity of **1** nor was cyclization of the pendant olefin in **34** detected.

Exchange of CO_2 via carbanion equivalents without competing quenching by other electrophiles (ketones, aldehydes, weak Brønsted acids) likely stems in part from the relatively high solubility of CO_2 in DMF and the slow kinetics of CO_2 evaporation into the reaction vessel headspace. For example, a 0.25 M solution of $^{13}\text{CO}_2$ in DMF retains a concentration of 0.2 M under a headspace of N_2 over one day (measured by ^{13}C NMR). For substrate **1** the rate of CO_2 exchange at 70°C was

~10-fold faster than for reaction with benzaldehyde. Perhaps counterintuitively, the rate of protodecarboxylation by weak Brønsted acids is inversely related to acidity (Fig. 3A, in order of decreasing rate of protonolysis: piperidine, aniline, methanol, phenol; see fig. S7 for details). This observation could be attributed to the relative capacity of these species to act as nucleophiles to sequester the liberated CO₂. Trapping of CO₂ prevents back reaction of the carbanion to the carboxylate leading to an increase in the observed rate of protodecarboxylation.

At 70°C under N₂, **1** underwent slow net carboxylate/proton metathesis to generate a half equivalent of the protodecarboxylated product **62** and a half equivalent of the CO₂-trapped malonate **63** (Fig. 3B). Product **62** likely arises from deprotonation of a second equivalent of aryl acetate to generate a dienolate nucleophile. The dienolate intermediate can react with the CO₂ released by the initial decarboxylation event. This observation demonstrates the striking efficiency of CO₂ capture by carbon nucleophiles under suitable conditions. Alkyl arenes generated by protodecarboxylation does not convert back to the carboxylate under the conditions where carboxylate exchange is observed (Fig. 3C). Carbonic anhydride intermediates are likely generated under the reaction conditions on the basis of the observed increase in α -carboxyl H/D exchange rates with **1-H₂** and **1-D₂** under CO₂ (Fig. 3D, see fig. S9 for details). The generation of a dienolate from the more acidic potassium carbonic anhydride may explain these reactivity differences. Direct detection of anhydride intermediates was not achieved. The capacity of non-enolizable carboxylates, such as **23** and **24**, to undergo reversible decarboxylation indicates that dienolate or enol intermediates are not essential for carboxylate exchange.

Finally, with an understanding of the factors that contribute to transient substrate decarboxylation, conditions were identified that allowed for the direct decarboxylative trapping of alternative classes of electrophiles (Fig. 3E). Carbon-carbon bond forming reactions by the trapping of aldehydes (**64–69**), a trifluoromethyl ketone (**70**), and an α,β -unsaturated ester (**71**) occurred in reasonable yields under conditions similar to those for reversible decarboxylation. The rates of product formation in aldehyde trapping experiments correlate with substrate electrophilicity. In some cases, it was beneficial (but not essential) to add 18-C-6 to improve decarboxylative reactivity or aniline to sequester liberated CO₂ (see SM for complete details). H/D-exchange followed by deutero-decarboxylation provides a simple approach to prepare CD₃ labeled toluenes and heterocycles from D₂O (**72–75**).

Efficient reversible decarboxylation/carboxylation masks the inherent reactivity of otherwise stable carboxylates. An appreciation of this phenomenon enables simple, direct protocols for isotopic exchange of carboxylic acids with ¹³CO₂ and methods for decarboxylative carbon-carbon bond forming

reactions. The potential for reversible decarboxylation should be considered more generally when designing and executing decarboxylative functionalization processes.

REFERENCES AND NOTES

1. T. Li, L. Huo, C. Pulley, A. Liu, Decarboxylation mechanisms in biological system. *Bioorg. Chem.* **43**, 2–14 (2012). [doi:10.1016/j.bioorg.2012.03.001](https://doi.org/10.1016/j.bioorg.2012.03.001) [Medline](#)
2. P. A. Frey, A. D. Hegeman, *Enzymatic Reaction Mechanisms* (Oxford Univ. Press, 2007).
3. M. H. O'Leary, in *The Enzymes*, D. S. Sigman, Ed. (Academic Press, 1992), vol. 20, pp. 235–269.
4. R. Kluger, Decarboxylation, CO₂ and the reversion problem. *Acc. Chem. Res.* **48**, 2843–2849 (2015). [doi:10.1021/acs.accounts.5b00306](https://doi.org/10.1021/acs.accounts.5b00306) [Medline](#)
5. A. J. Colussi, V. T. Amorebieta, M. A. Grela, Very low pressure pyrolysis of phenylacetic acid. *J. Chem. Soc., Faraday Trans.* **88**, 2125–2127 (1992). [doi:10.1039/f9928802125](https://doi.org/10.1039/f9928802125)
6. N. Rodríguez, L. J. Goossen, Decarboxylative coupling reactions: A modern strategy for C–C-bond formation. *Chem. Soc. Rev.* **40**, 5030–5048 (2011). [doi:10.1039/c1cs15093f](https://doi.org/10.1039/c1cs15093f) [Medline](#)
7. A. K. Vijh, B. E. Conway, Electrode kinetic aspects of the Kolbe reaction. *Chem. Rev.* **67**, 623–664 (1967). [doi:10.1021/cr60250a003](https://doi.org/10.1021/cr60250a003)
8. Z. W. Zuo, D. T. Ahneman, L. Chu, J. A. Terrett, A. G. Doyle, D. W. C. MacMillan, Merging photoredox with nickel catalysis: Coupling of α -carboxyl sp³-carbons with aryl halides. *Science* **345**, 437–440 (2014). [doi:10.1126/science.1255525](https://doi.org/10.1126/science.1255525) [Medline](#)
9. D. H. R. Barton, S. D. Gero, B. Quiclet-Sire, M. Samadi, N. Ozbalik, J. C. Sarma, M. Ramesh, New reactions for use in natural products chemistry. *Pure Appl. Chem.* **60**, 1549–1554 (1988). [doi:10.1351/pac19886011549](https://doi.org/10.1351/pac19886011549)
10. T. Qin, J. Cornella, C. Li, L. R. Malins, J. T. Edwards, S. Kawamura, B. D. Maxwell, M. D. Eastgate, P. S. Baran, A general alkyl-alkyl cross-coupling enabled by redox-active esters and alkylzinc reagents. *Science* **352**, 801–805 (2016). [doi:10.1126/science.aaf6123](https://doi.org/10.1126/science.aaf6123) [Medline](#)
11. M. B. Smith, *March's Advanced Organic Chemistry – Reactions, Mechanisms, and Structure* (Wiley, ed. 7, 2013).
12. J. Schwarz, B. König, Decarboxylative reactions with and without light – a comparison. *Green Chem.* **20**, 323–361 (2018). [doi:10.1039/C7GC02949G](https://doi.org/10.1039/C7GC02949G)
13. R. B. Silverman, Ed., *Organic Chemistry of Enzyme-Catalyzed Reactions* (Academic Press, ed. 2, 2002), pp. 289–320.
14. S. S. Bailey, K. A. P. Payne, A. Saaret, S. A. Marshall, I. Gostimskaya, I. Kosov, K. Fisher, S. Hay, D. Leys, Enzymatic control of cycloadduct conformation ensures reversible 1,3-dipolar cycloaddition in a prFMN-dependent decarboxylase. *Nat. Chem.* **11**, 1049–1057 (2019). [doi:10.1038/s41557-019-0324-8](https://doi.org/10.1038/s41557-019-0324-8) [Medline](#)
15. T. Nunoura, Y. Chikaraishi, R. Izaki, T. Suwa, T. Sato, T. Harada, K. Mori, Y. Kato, M. Miyazaki, S. Shimamura, K. Yanagawa, A. Shuto, N. Ohkouchi, N. Fujita, Y. Takaki, H. Atomi, K. Takai, A primordial and reversible CO₂ cycle in a facultatively chemolithoautotrophic thermophile. *Science* **359**, 559–563 (2018). [doi:10.1126/science.aao3407](https://doi.org/10.1126/science.aao3407) [Medline](#)
16. A. Mall, J. Sobotta, C. Huber, C. Tschirner, S. Kowarschik, K. Bačnik, M. Mergelsberg, M. Boll, M. Hügler, W. Eisenreich, I. A. Berg, Reversibility of citrate synthase allows autotrophic growth of a thermophilic bacterium. *Science* **359**, 563–567 (2018). [doi:10.1126/science.aao2410](https://doi.org/10.1126/science.aao2410) [Medline](#)
17. X. Sheng, Y. Patskovsky, A. Vladimirova, J. B. Bonanno, S. C. Almo, F. Himo, F. M. Raushel, Mechanism and structure of γ -resorcyate decarboxylase. *Biochemistry* **57**, 3167–3175 (2018). [doi:10.1021/acs.biochem.7b01213](https://doi.org/10.1021/acs.biochem.7b01213) [Medline](#)
18. Q. Liu, L. Wu, R. Jackstell, M. Beller, Using carbon dioxide as a building block in organic synthesis. *Nat. Commun.* **6**, 5933 (2015). [doi:10.1038/ncomms6933](https://doi.org/10.1038/ncomms6933) [Medline](#)
19. A. Tortajada, F. Juliá-Hernández, M. Börjesson, T. Moragas, R. Martin, Transition-metal-catalyzed carboxylation reactions with carbon dioxide. *Angew. Chem. Int. Ed.* **57**, 15948–15982 (2018). [doi:10.1002/anie.201803186](https://doi.org/10.1002/anie.201803186) [Medline](#)
20. D. J. Darensbourg, E. M. Longridge, M. W. Holtcamp, K. K. Klausmeyer, J. H. Reibenspies, Reversible decarboxylation of phosphine derivatives of Cu(I) cyanoacetate. Mechanistic aspects germane to catalytic decarboxylation of carboxylic acids. *J. Am. Chem. Soc.* **115**, 8839–8840 (1993). [doi:10.1021/ja00072a047](https://doi.org/10.1021/ja00072a047)
21. A. Häussermann, F. Rominger, B. F. Straub, CO₂ on a tightrope: Stabilization, room-

- temperature decarboxylation, and sodium-induced carboxylate migration. *Chemistry* **18**, 14174–14185 (2012). [doi:10.1002/chem.201202298](https://doi.org/10.1002/chem.201202298) [Medline](#)
22. A. Szabolcs, J. Szammer, L. Noszkó, A new method for the preparation of carboxyl-labelled aliphatic carboxylic acids. *Tetrahedron* **30**, 3647–3648 (1974). [doi:10.1016/S0040-4020\(01\)97049-8](https://doi.org/10.1016/S0040-4020(01)97049-8)
 23. R. Nakai, M. Sugii, H. Nakao, Isotopic tracer studies of the ketonic pyrolysis of sodium carboxylates. *J. Am. Chem. Soc.* **81**, 1003–1006 (1959). [doi:10.1021/ja01513a061](https://doi.org/10.1021/ja01513a061)
 24. P. J. Moon, S. Yin, R. J. Lundgren, Ambient decarboxylative arylation of malonate half-esters via oxidative catalysis. *J. Am. Chem. Soc.* **138**, 13826–13829 (2016). [doi:10.1021/jacs.6b08906](https://doi.org/10.1021/jacs.6b08906) [Medline](#)
 25. P. J. Moon, A. Fahandj-Sadi, W. Qian, R. J. Lundgren, Decarboxylative benzylation of aryl and alkenyl boronic esters. *Angew. Chem. Int. Ed.* **57**, 4612–4616 (2018). [doi:10.1002/anie.201800829](https://doi.org/10.1002/anie.201800829) [Medline](#)
 26. G. Destro, K. Horkka, O. Loreau, D. A. Buisson, L. Kingston, A. Del Vecchio, M. Schou, C. S. Elmore, F. Taran, T. Cantat, D. Audisio, Transition-metal-free carbon isotope exchange of phenyl acetic acids. *Angew. Chem. Int. Ed.* (2020). [doi:10.1002/anie.202002341](https://doi.org/10.1002/anie.202002341) [Medline](#)
 27. D. H. Hunter, M. Hamity, V. Patel, R. A. Perry, Crown ether catalysis of decarboxylation: A general survey of reactivity and detailed analysis of the triphenylacetate anion. *Can. J. Chem.* **56**, 104–113 (1978). [doi:10.1139/v78-017](https://doi.org/10.1139/v78-017)
 28. C. S. Elmore, R. A. Bragg, Isotope chemistry: a useful tool in the drug discovery arsenal. *Bioorg. Med. Chem. Lett.* **25**, 167–171 (2015). [doi:10.1016/j.bmcl.2014.11.051](https://doi.org/10.1016/j.bmcl.2014.11.051) [Medline](#)
 29. K. Hinsinger, G. Pieters, The emergence of carbon isotope exchange. *Angew. Chem. Int. Ed.* **58**, 9678–9680 (2019). [doi:10.1002/anie.201905368](https://doi.org/10.1002/anie.201905368) [Medline](#)
 30. C. Kingston, M. A. Wallace, A. J. Allentoff, J. N. deGruyter, J. S. Chen, S. X. Gong, S. Bonacorsi Jr., P. S. Baran, Direct carbon isotope exchange through decarboxylative carboxylation. *J. Am. Chem. Soc.* **141**, 774–779 (2019). [doi:10.1021/jacs.8b12035](https://doi.org/10.1021/jacs.8b12035) [Medline](#)
 31. A. Tortajada, Y. Duan, B. Sahoo, F. Cong, G. Toupalas, A. Sallustrau, O. Loreau, D. Audisio, R. Martin, Catalytic decarboxylation/carboxylation platform for accessing isotopically labeled carboxylic acids. *ACS Catal.* **9**, 5897–5901 (2019). [doi:10.1021/acscatal.9b01921](https://doi.org/10.1021/acscatal.9b01921)
 32. R. Voges, J. R. Heyes, T. Moenius, *Preparation of Compounds Labeled with Tritium and Carbon-14* (Wiley, 2009).
 33. D. U. Nielsen, K. T. Neumann, A. T. Lindhardt, T. Skrydstrup, Recent developments in carbonylation chemistry using [¹³C]CO, [¹³C]CO, and [¹⁴C]CO. *J. Labelled Comp. Radiopharm.* **61**, 949–987 (2018). [doi:10.1002/jlcr.3645](https://doi.org/10.1002/jlcr.3645) [Medline](#)
 34. D. R. Gauthier Jr., N. R. Rivera, H. Yang, D. M. Schultz, C. S. Shultz, Palladium-catalyzed carbon isotope exchange on aliphatic and benzoic acid chlorides. *J. Am. Chem. Soc.* **140**, 15596–15600 (2018). [doi:10.1021/jacs.8b09808](https://doi.org/10.1021/jacs.8b09808) [Medline](#)
 35. G. Destro, O. Loreau, E. Marcon, F. Taran, T. Cantat, D. Audisio, Dynamic carbon isotope exchange of pharmaceuticals with labeled CO₂. *J. Am. Chem. Soc.* **141**, 780–784 (2019). [doi:10.1021/jacs.8b12140](https://doi.org/10.1021/jacs.8b12140) [Medline](#)
 36. I. Shiina, K. Ono, K. Nakata, Non-enzymatic dynamic kinetic resolution of racemic α-arylalkanoic acids: An advanced asymmetric synthesis of chiral nonsteroidal anti-inflammatory drugs (NSAIDs). *Catal. Sci. Technol.* **2**, 2200–2205 (2012). [doi:10.1039/c2cy20329d](https://doi.org/10.1039/c2cy20329d)
 37. P. J. Harrington, E. Lodewijk, Twenty years of naproxen technology. *Org. Process Res. Dev.* **1**, 72–76 (1997). [doi:10.1021/op960009e](https://doi.org/10.1021/op960009e)
 38. T. Ooi, K. Maruoka, Recent advances in asymmetric phase-transfer catalysis. *Angew. Chem. Int. Ed.* **46**, 4222–4266 (2007). [doi:10.1002/anie.200601737](https://doi.org/10.1002/anie.200601737) [Medline](#)
 39. W. C. Still, M. Kahn, A. Mitra, Rapid chromatographic technique for preparative separations with moderate resolution. *J. Org. Chem.* **43**, 2923–2925 (1978). [doi:10.1021/jo00408a041](https://doi.org/10.1021/jo00408a041)
 40. R. Shang, Z.-W. Yang, Y. Wang, S.-L. Zhang, L. Liu, Palladium-catalyzed decarboxylative couplings of 2-(2-azaaryl)acetates with aryl halides and triflates. *J. Am. Chem. Soc.* **132**, 14391–14393 (2010). [doi:10.1021/ja107103b](https://doi.org/10.1021/ja107103b) [Medline](#)
 41. J. E. Day, S. Y. Sharp, M. G. Rowlands, W. Aherne, P. Workman, C. J. Moody, Targeting the Hsp90 chaperone: Synthesis of novel resorcylic acid macrolactone inhibitors of Hsp90. *Chemistry* **16**, 2758–2763 (2010). [doi:10.1002/chem.200902766](https://doi.org/10.1002/chem.200902766) [Medline](#)
 42. R. Shang, Z. Huang, L. Chu, Y. Fu, L. Liu, Palladium-catalyzed decarboxylative coupling of potassium nitrophenyl acetates with aryl halides. *Org. Lett.* **13**, 4240–4243 (2011). [doi:10.1021/ol201750s](https://doi.org/10.1021/ol201750s) [Medline](#)
 43. T. T. Mai, B. Viswambaran, D. Gori, R. Guillot, J.-V. Naubron, C. Kouklovsky, V. Alezra, Frozen chirality of tertiary aromatic amides: Access to enantioenriched tertiary α-amino acid or amino alcohol without chiral reagent. *Chemistry* **23**, 5787–5798 (2017). [doi:10.1002/chem.201700110](https://doi.org/10.1002/chem.201700110) [Medline](#)
 44. P. Borowiecki, P. Wińska, M. Bretner, M. Gizińska, M. Koroniewicz, M. Staniszewska, Synthesis of novel proxiphylline derivatives with dual anti-*Candida albicans* and anticancer activity. *Eur. J. Med. Chem.* **150**, 307–333 (2018). [doi:10.1016/j.ejmech.2018.02.077](https://doi.org/10.1016/j.ejmech.2018.02.077) [Medline](#)
 45. D. J. D. Bon, O. Kováč, V. Ferugová, F. Zálešák, J. Pospíšil, One and two-carbon homologation of primary and secondary alcohols to corresponding carboxylic esters using β-carbonyl BT sulfones as a common intermediate. *J. Org. Chem.* **83**, 4990–5001 (2018). [doi:10.1021/acs.joc.8b00112](https://doi.org/10.1021/acs.joc.8b00112) [Medline](#)
 46. R. Shintani, S. Park, F. Shirozu, M. Murakami, T. Hayashi, Palladium-catalyzed asymmetric decarboxylative lactamization of γ-methylidene-δ-valerolactones with isocyanates: Conversion of racemic lactones to enantioenriched lactams. *J. Am. Chem. Soc.* **130**, 16174–16175 (2008). [doi:10.1021/ja807662b](https://doi.org/10.1021/ja807662b) [Medline](#)
 47. Y. Li, W. Zhang, H. Zhang, W. Tian, L. Wu, S. Wang, M. Zheng, J. Zhang, C. Sun, Z. Deng, Y. Sun, X. Qu, J. Zhou, Structural basis of a broadly selective acyltransferase from the polyketide synthase of splenocin. *Angew. Chem. Int. Ed.* **57**, 5823–5827 (2018). [doi:10.1002/anie.201802805](https://doi.org/10.1002/anie.201802805) [Medline](#)
 48. J. Nandi, N. E. Leadbeater, Visible-light-driven catalytic oxidation of aldehydes and alcohols to nitriles by 4-acetamido-TEMPO using ammonium carbamate as a nitrogen source. *Org. Biomol. Chem.* **17**, 9182–9186 (2019). [doi:10.1039/C9OB01918A](https://doi.org/10.1039/C9OB01918A) [Medline](#)
 49. T. Kimachi, H. Nagata, Y. Kobayashi, K. Takahashi, E. Torii, M. Ju-ichi, Regioselective ring-opening reaction of unsymmetrical 2,3-diaryl epoxides via catalytic hydrogenolysis with Pd(0)EnCat™. *Synlett* **2011**, 365–368 (2011). [doi:10.1055/s-0030-1259505](https://doi.org/10.1055/s-0030-1259505)

ACKNOWLEDGMENTS

We thank A. Speed for analysis of this manuscript and experimental suggestions. We thank J. Takats for the gift of ¹³CO₂ to initiate this project and B. Reiz for assistance in mass spectrometry data analysis. **Funding:** Support was provided by NSERC Canada (RGPIN-2019-06050, RGPAS-2019-00051) the Canadian Foundation for Innovation (IOF 32691), the University of Alberta, and the Killam Trusts **Author contributions:** P.J.M. and R.J.L. conceived the study. Experiments and data analysis were conducted by D.K., P.J.M., E.J.K.L. and O.B. The manuscript was written by R.J.L. with consultation from all co-authors. **Competing interests:** P.J.M. and R.J.L. (University of Alberta) are pursuing a provisional patent on this work. **Data and materials availability:** All experimental data are available in the supplementary materials.

SUPPLEMENTARY MATERIALS

science.sciencemag.org/cgi/content/full/science.abb4129/DC1

Materials and Methods

Figs. S1 to S9

HRMS and NMR Spectra

References (39–49)

25 February 2020; accepted 9 June 2020

Published online 18 June 2020

10.1126/science.abb4129

A enzymatic decarboxylation – facile under physiological conditions

B unactivated carboxylic acids construed to need high temperature to decarboxylate

C

D

cation	¹³ C incorp. (time)
K ⁺	85% (1 h)
H ⁺	<5% (8 h)
Li ⁺	72% (8 h)
Na ⁺	77% (8 h)
Cs ⁺	63% (5 h, rt)
0.5 Zn ²⁺ /Cu ²⁺	<5% (8 h)

conditions	¹³ C incorp. (time)
THF	0% (5 h)
DCE	0% (5 h)
H ₂ O	0% (5 h)
DMSO	47% (5 h)
DMA	45% (5 h)
DMF/18-C-6	67% (5 h)

Fig. 1. Overview of decarboxylative processes and carboxylate exchange. (A) Decarboxylation catalyzed by enzymes under physiological conditions (PDB ID: 2INF). (B) Comparison of conditions used for thermal decarboxylation of different substrates. (C) Comparison of CO₂ exchange (red) and protonation with MeOH (black) for 4-cyanophenylacetate (**1**). (D) Impact of salt and reaction conditions. E⁺, electrophile; DMF, dimethylformamide; THF, tetrahydrofuran; DCE, 1,2-dichloroethane; DMSO, dimethylsulfoxide; DMA, dimethylacetamide; 18-C-6, 18-crown-6.

Fig. 2. Carboxylate exchange scope and application. Unless noted yields are of isolated material. *Calibrated ¹H NMR spectroscopy yield. †1 Equivalent 18-C-6 added. ‡%¹³C Incorporation and yield determined by analysis of the corresponding methyl or benzyl ester. § DMSO used instead of DMF. See the supplementary materials for complete details. NMR, nuclear magnetic resonance.

Fig. 3. Mechanistic control experiments and electrophile trapping. (A) Relative rates for arylacetic acid salt protodecarboxylation by Brønsted acids and decarboxylative trapping by benzaldehyde. (B) Net carboxylate/proton metathesis of **1** in the absence of additional CO₂. (C) CO₂ transfer between arylacetic acid salt and alkylarene does not occur. (D) The presence of CO₂ accelerates the rate of methylene C–H/D exchange in **1**. (E) Scope examples for the decarboxylative trapping of alternative classes of electrophiles. See the supplementary materials for complete details and full reaction conditions. *Yield determined by ¹H NMR spectroscopy. Ar, (4-CN)C₆H₄.